

The Biological Perspective

2

▲ TABLE OF CONTENTS

To access the resource listed, click on the hot linked title or press **CTRL + click**

To return to the Table of Contents, click on **▲Return to Table of Contents**

To return to a section of the Lecture Guide, click on **►Return to Lecture Guide**

► LECTURE GUIDE

- Neurons and Nerves: Building the Network (p. 3)
- An Overview of the Nervous System (p. 4)
- Distant Connections: The Endocrine System (p. 4)
- Looking Inside the Living Brain (p. 5)
- From the Bottom Up: Structures of the Brain (p. 5)
- Applying Psychology to Everyday Life: Paying Attention to Attention-Deficit/Hyperactivity Disorder (p. 7)
- Chapter Summary (p. 7)

► FULL CHAPTER RESOURCES

- Chapter-At-A-Glance (p. 2)
- Lecture Guide (p. 3)
- Learning Objectives (p. 8)
- Rapid Review (p. 9)
- Changes from the Fourth Edition to the Fifth Edition (p. 10)
- Lecture Launchers and Discussion Topics (p. 11)
- Classroom Activities, Demonstrations, and Exercises (p. 26)
- Handout Masters (p. 36)
- Video Series (p. 44)
- Visual Brain (p. 45)
- Experiment and Survey Simulations (p. 46)
- Writing Space and Writing Practice (p. 47)

Chapter 2 Answer Key

Practice Quiz 2.1

1. a; 2. b; 3. b; 4. a; 5. a; 6. d

Practice Quiz 2.2

1. a; 2. d; 3. b; 4. b; 5. a

Practice Quiz 2.3

Copyright © 2017 Pearson Education. All rights reserved.

2-1

1. a; 2. c; 3. b; 4. c

Practice Quiz 2.4

1. d; 2. d; 3. a; 4. b

Practice Quiz 2.5

1. c; 2. a; 3. d; 4. d; 5. c; 6. a

Test Yourself 2

1. b; 2. c; 3. c; 4. a; 5. a; 6. c; 7. b; 8. b; 9. d; 10. b; 11. b; 12. d; 13. c;
14. d; 15. a; 16. a; 17. d; 18. b; 19. d; 20. b

▼ CHAPTER-AT-A-GLANCE

<i>Detailed Outline</i>	<i>Instructor Resources</i>	<i>Multimedia Resources</i>
Neurons and Nerves: Building the Network Structure of the Neuron Generating the Message Sending the Message Neurotransmission	Learning Objectives: 2.1, 2.2, 2.3 Lecture Launchers: 2.1, 2.2, 2.3, 2.4 Activities & Exercises: 2.1, 2.2, 2.3	Video: <i>The Basics: How the Brain Works, Part 1</i> (5:04) — Find out about neurons: their parts, how they communicate, and their function in the brain. Video: <i>In the Real World: Neurotransmitters</i> (3:10) — Learn about the neurotransmitter dopamine and what happens when there is an imbalance of dopamine as a result of natural causes or drugs.
An Overview of the Nervous System The Central Nervous System The Peripheral Nervous System	Learning Objectives: 2.4, 2.5 Lecture Launchers: 2.5, 2.6, 2.7 Activities & Exercises: 2.4, 2.5, 2.6	Video: <i>The Basics: How the Brain Works, Part 2</i> (6:11) — Find out about the nervous system: how it's divided and processes information, the significance of each brain structure, and how neuronal transmission works. Video: <i>Special Topics: The Plastic Brain</i> (7:04) — Learn specifics about how the human brain grows and develops throughout the lifespan and how the brain can remarkably repair itself or compensate after being damaged. Video: <i>The Big Picture: My Brain Made Me Do It</i> (4:59) — Learn about the function of different parts of the nervous system and how it integrates multiple signals to create experience. Video: <i>What's In It For Me? Your Brain on Drugs</i> (3:51) — Neurotransmitter imbalances can occur from chemicals that we willingly ingest, such as drugs or alcohol.
Distant Connections: The Endocrine Glands The Pituitary Other Endocrine Glands	Learning Objectives: 2.6, 2.7 Lecture Launchers: 2.7	
Looking Inside the Living Brain Methods for Studying Specific Brain Regions Neuroimaging Techniques	Learning Objectives: 2.8, 2.9 Lecture Launchers: 2.8, 2.9, 2.10, 2.11, 2.12	
Structures of the Brain The Hindbrain The Limbic System The Cortex Association Areas of the Cortex Cerebral Hemispheres	Learning Objectives: 2.10, 2.11, 2.12, 2.13, 2.14 Lecture Launchers: 2.13, 2.14, 2.15, 2.16, 2.17, 2.18, 2.19, 2.20 Activities & Exercises: 2.7, 2.8, 2.9, 2.10 Handout: 2.1	Video: <i>Thinking Like a Psychologist: The Pre-Frontal Cortex</i> (3:28) — See how certain areas of the brain contribute to different behaviors, and how both biological and environmental factors can lead a person to become a violent criminal later in life. Writing Assignment: Your best friends invite you over for pizza and a friendly game of cards. Describe how the following parts of the brain are involved during your evening of eating pizza, socializing, and playing cards: Broca's area, hippocampus, hypothalamus, and occipital lobe.
Applying Psychology to Everyday Life: Paying Attention to ADHD	Learning Objective: 2.15	

▼ Lecture Guide

I. NEURONS AND NERVES: BUILDING THE NETWORK

Lecture Launchers and Discussion Topics

- 2.1 Neurotransmitters: Chemical Communicators of the Nervous System
 - 2.2 Exceptions to the Rules
 - 2.3 The Glue of Life: Neuroglial Cells
 - 2.4 My Mother, Myself

Classroom Activities, Demonstrations, and Exercises

- 2.1 - Using Dominoes to Understand the Action Potential
 - 2.2 - Stemming the Tide of Misinformation
 - 2.3 - Environmental Influences on the Brain
- Video - In the Real World: Neurotransmitters
- Video - Basics: How the Brain Works, Part 1
- Animation - The Nerve Impulse in Afferent and Efferent Neurons
 - Animation - Structure of the Neuron
 - Animation - Neural Impulse Action Potential
 - Animation - The Synapse
 - Animation - Neurotransmitters: Reuptake
 - Audio - Endocrine System (Podcast)

Learning Objective 2.1 - Identify the parts of a neuron and the function of each.

- A. Structure of the neuron: The nervous system's building block
 - 1. Neurons: Dendrites, soma, axons
 - a. Myelin insulates and protects axons, speeds neural conduction
 - 2. Glial cells support, separate, and insulate neurons from each other

Learning Objective 2.2 - Explain the action potential.

- B. Generating the message within the neuron: The neural impulse
 - 1. At rest, the neuron is negatively charged inside and positively charged outside
 - 2. When stimulated, the charge reverses
 - a. This is the action potential
 - 3. Neurons fire in an all-or-nothing manner

Learning Objective 2.3 - Describe how neurons use neurotransmitters to communicate with each other and with the body.

- C. Sending the message to other cells: The synapse
 - 1. Synaptic vesicles release neurotransmitter into the synapse
 - 2. Neurotransmitter molecules fit into receptor sites
 - a. Neurotransmitters may be either excitatory or inhibitory
- D. Neurotransmitters: Messengers of the network
 - 1. Acetylcholine
 - 2. GABA
 - 3. Serotonin
 - 4. Dopamine
 - 5. Endorphins
- E. Cleaning up the synapse: Reuptake and enzymes
 - 1. Most neurotransmitters are taken back into the synaptic vesicles
 - 2. Acetylcholine is cleared out of the synapse by enzymes that break up the molecules

Practice Quiz 2.1 Answer Key

1. a; 2. b; 3. b; 4. a; 5. a; 6. d

▲ Return to Chapter 2: Table of Contents

II. AN OVERVIEW OF THE NERVOUS SYSTEM

Lecture Launchers and Discussion Topics

- 2.5 Brain Metaphors
- 2.6 The Cranial Nerves
- 2.7 Hormone Imbalances

Classroom Activities, Demonstrations, and Exercises

- 2.4 - Demonstrating Neural Conduction: The Class as a Neural Network
 - 2.5 - The Dollar Bill Drop
- 2.6 - Reaction Time and Neural Processing

○ Video - Basics: How the Brain Works, Part 2

○ Video - The Big Picture: My Brain Made Me Do It

○ Video - What's In It For Me? Your Brain On Drugs

○ Animation - The Spinal Cord Reflex

○ Simulation - The Nerve Impulse in Afferent and Efferent Neurons

Learning Objective 2.4 - Describe how the brain and spinal cord interact and respond to external experiences.

A. The central nervous system

1. Brain
2. Spinal cord
3. Neuroplasticity and neurogenesis offer exciting possibilities for cell

regeneration

Learning Objective 2.5 - Differentiate the roles of the somatic and autonomic nervous systems.

B. The peripheral nervous system

1. The somatic nervous system
 - a. Sensory pathway
 - b. Motor pathway
2. The autonomic nervous system
 - a. Sympathetic division
 - b. Parasympathetic division

Practice Quiz 2.2 Answer Key

1. a; 2. d; 3. b; 4. b; 5. a

III. DISTANT CONNECTIONS: THE ENDOCRINE GLANDS

Lecture Launchers and Discussion Topics

- 2.8 Hormone Imbalances

Learning Objective 2.6 - Explain why the pituitary gland is known as the "master gland."

- A. Pituitary gland
 - 1. Just below the hypothalamus
 - 2. Controls the levels of salt and water in the system
 - 3. In women, controls the onset of labor and lactation
 - a. "Love hormones" and "trust hormones" have popular appeal

Learning Objective 2.7 - Recall the role of various endocrine glands.

- B. Other endocrine glands
 - 1. Pineal gland: secretes melatonin
 - 2. Thyroid gland: controls metabolism by secreting thyroxin
 - 3. Pancreas: controls sugar in the blood by secreting insulin and glucagons
 - 4. Gonads: ovaries and testes regulate sexual growth, activity, reproduction
 - 5. Adrenal glands: on top of each kidney; control stress reactions

Practice Quiz 2.3 Answer Key

1. a; 2. c; 3. b; 4. c

▲ Return to Chapter 2: Table of Contents

IV. LOOKING INSIDE THE LIVING BRAIN

Lecture Launchers and Discussion Topics

- 2.9 Psychophysiological Measurement
 - 2.10 Berger's Wave
 - 2.11 Lie Detectors 2.0
 - 2.12 Women, Men, and PETs
- 2.13 Using fMRI and MEG to Study Phantom Limb Pain

○ Video - Brain Imaging

Learning Objective 2.8 - Describe how lesioning studies and brain stimulation are used to study the brain.

- A. Lesioning studies
 - 1. Destroy certain areas of the brain in animals
 - 2. Case studies of human brain damage
- B. Brain stimulation
 - 1. Deep brain stimulation (DBS) is an invasive technique
 - a. Optogenetics may offer a comparable alternative
 - 2. TMS, rTMS, and tDCS are noninvasive methods for stimulating the brain

Learning Objective 2.9 - Compare and contrast neuroimaging techniques for mapping the structure and function of the brain.

- C. Mapping structure
 - 1. CT scans are computer-aided X-rays that show the skull and brain structure

2. MRI scans use a magnetic field, radio pulses, and a computer
3. Diffusion tensor imaging (DTI) off to a promising start
- D. Mapping function
 1. EEG uses electrodes placed on the scalp that are then amplified and viewed
 2. ERPs offers a glimpse of activity in response to a stimulus
 3. MEG directly identifies areas of brain activation
 4. PET scans use a radioactive sugar injected into the bloodstream
 5. fMRI allows researchers to look at the activity of the brain over a time period.

Practice Quiz 2.4 Answer Key

1. d; 2. d; 3. a; 4. b

▲ Return to Chapter 2: Table of Contents

V. FROM THE BOTTOM UP: THE STRUCTURES OF THE BRAIN

Lecture Launchers and Discussion Topics

- 2.14 - The Importance of a Wrinkled Cortex
 - 2.15 - Brain's Bilingual Broca
 - 2.16 - A New Look at Phineas Gage
 - 2.17 - Freak Accidents and Brain Injuries
 - 2.18 - Understanding Hemispheric Function
- 2.19 - Handedness, Eyedness, Footedness, Facedness
 - 2.20 - Workplace Problems: Left-Handedness
 - 2.21 - The Results of a Hemispherectomy

Classroom Activities, Demonstrations, and Exercises

- 2.7 - Mapping the Brain
- 2.8 - Football and Brain Damage
- 2.9 - Hemispheric Lateralization
- 2.10 - Hemispheric Communication and the Split Brain

○ Video - Thinking Like a Psychologist: The Prefrontal Cortex

○ Video - Special Topics: The Plastic Brain

○ Animation - The Limbic System

○ Animation - The Split-Brain Experiment

○ Simulation - Hemispheric Specialization

Learning Objective 2.10 - Identify the different structures of the hindbrain and the function of each.

A. The hindbrain

1. *Pons*—involved in sleeping, waking, and dreaming
2. *Medulla*—regulates breathing and heart rate; automatic functions
3. *Reticular formation*—responsible for arousal and alertness
4. *Cerebellum*—regulates balance and coordination of movement

Learning Objective 2.11 - Identify the structures of the brain that are involved in emotion, learning, memory, and motivation.

B. Structures under the cortex

1. *Thalamus*—directs incoming sensory messages to higher centers
2. *Hypothalamus*—associated with drives, such as hunger, thirst, emotion, sex

3. *Hippocampus*—"gateway to memory" that enables us to form new memories
4. *Amygdala*—evaluates sensory information to determine its importance
5. *Cingulate cortex*—regions contribute to different aspects of emotion and cognition

Learning Objective 2.12 - Identify the parts of the cortex that process the different senses and those that control movement of the body.

C. Cortex

1. Connected by band of fibers called corpus callosum
2. Right hemisphere in charge of left side of the body
3. Left hemisphere in charge of right side of the body
4. Lateralization—each hemisphere has somewhat different tasks and talents
5. Divided into four regions
 - a. *Occipital lobes*—contain the visual cortex
 - b. *Parietal lobes*—contain somatosensory cortex, that receives information about pressure, pain, touch, and temperature from all over the body
 - c. *Temporal lobes*—contain auditory cortex
 - d. *Frontal lobes*—contain the motor cortex; responsible for making plans, taking initiative, and thinking creatively

Learning Objective 2.13 - Name the parts of the cortex that are responsible for higher forms of thought, such as language.

D. Association areas of the cortex

1. Association areas are found in all the lobes but particularly in the frontal lobes
2. Help make sense of information received from lower areas of the brain
3. Broca's area is responsible for producing fluent, understandable speech
4. Wernicke's area is responsible for the understanding of language

Learning Objective 2.14 - Explain how some brain functions differ between the left and right hemispheres.

E. The cerebral hemispheres: Are you in your right mind?

1. Left side of the brain seems to control language, writing, logical thought, analysis, and mathematical abilities
2. The right side of the brain processes information globally and controls emotional expression, spatial perception, and recognition of faces, patterns, melodies, and emotions
3. Information presented only to the left hemisphere can be verbalized but information only sent to the right cannot
 - a. Split-brain research, handedness, provide evidence of these properties

Practice Quiz 2.5 Answer Key

1. c; 2. a; 3. d; 4. d; 5. c; 6. a

▲ Return to Chapter 2: Table of Contents

VI. APPLYING PSYCHOLOGY TO EVERYDAY LIFE: PAYING ATTENTION TO ATTENTION-DEFICIT/HYPERACTIVITY DISORDER

Copyright © 2017 Pearson Education. All rights reserved.

- Video - Attention Deficit Hyperactivity Disorder (ADHD): Dr. Raun Mel

Learning Objective 2.15 - Identify some potential causes of attention-deficit/hyperactivity disorder.

- A. ADHD may be caused by the interaction of a variety of factors
- B. Often persists into adulthood

▲ Return to Chapter 2: Table of Contents

VI. CHAPTER SUMMARY

Classroom Activities, Demonstrations, and Exercises

- Crossword Puzzle
- Fill-in-the-Blanks

Test Yourself 2 Answer Key

1. b; 2. c; 3. c; 4. a; 5. a; 6. c; 7. b; 8. b; 9. d; 10. b; 11. b; 12. d; 13. c;
14. d; 15. a; 16. a; 17. d; 18. b; 19. d; 20. b

▲ Return to Chapter 2: Table of Contents

▼ LEARNING OBJECTIVES

- 2.1 Identify the parts of a neuron and the function of each.
- 2.2 Explain the action potential.
- 2.3 Describe how neurons use neurotransmitters to communicate with each other and with the body.
- 2.4 Describe how the brain and spinal cord interact and respond to external experiences.
- 2.5 Differentiate the roles of the somatic and autonomic nervous systems.
- 2.6 Explain why the pituitary gland is known as the "master gland."
- 2.7 Recall the role of various endocrine glands.
- 2.8 Describe how lesioning studies and brain stimulation are used to study the brain.
- 2.9 Compare and contrast neuroimaging techniques for mapping the structure and function of the brain.
- 2.10 Identify the different structures of the hindbrain and the function of each.
- 2.11 Identify the structures of the brain that are involved in emotion, learning, memory, and motivation.
- 2.12 Identify the parts of the cortex that process the different senses and those that control movement of the body.
- 2.13 Name the parts of the cortex that are responsible for higher forms of thought, such as language.
- 2.14 Explain how some brain functions differ between the left and right hemispheres.
- 2.15 Identify some potential causes of attention-deficit/hyperactivity disorder.

▲ Return to Chapter 2: Table of Contents

▼ RAPID REVIEW

The **nervous system** is composed of a complex network of cells throughout the body. The cells in the nervous system that carry information are called **neurons**. Information enters at the **dendrites**, flows through the cell body (or **soma**) and down the **axon**. Although neurons are the cells that carry the information, most of the nervous system consists of **glial cells** that provide food, support, and insulation to the neuron cells. The insulation around the neuron is called **myelin** and works in a way similar to the plastic coating on an electrical wire. Bundles of myelin-coated axons are wrapped together in cable-like structures called **nerves**. The movement of an electrical signal across a neuron is called an **action potential**. A neuron fires in an **all-or-none** manner: The neuron either has an action potential or it does not. When the electrical signal travels down the axon and reaches the other end of the neuron called the **axon terminal**, it causes the **neurotransmitters** in the **synaptic vesicles** to be released into the fluid-filled **synapse** between the two cells.

The **central nervous system (CNS)** is made up of the brain and the **spinal cord**. **Afferent (sensory) neurons** send information from the senses to the spinal cord, whereas **efferent (motor) neurons** send commands from the spinal cord to the muscles. **Interneurons** connect sensory and motor neurons and help to coordinate the signals. The **peripheral nervous system (PNS)** is made up of all the nerves and neurons that are NOT in the brain or spinal cord, and is divided into two parts: the **somatic nervous system** and the **autonomic nervous system**. The autonomic nervous system is in turn divided into two parts: the **sympathetic division** and the **parasympathetic division**.

The **endocrine glands** represent a second communication system in the body. The endocrine glands secrete chemicals called **hormones** directly into the bloodstream. The **pituitary gland** is located in the brain and secretes the hormones that control milk production, salt levels, and the activity of other glands. The **pineal gland** is also located in the brain and regulates the sleep cycle through the secretion of melatonin. The **thyroid gland** is located in the neck and releases a hormone that regulates metabolism. The **pancreas** controls the level of blood sugar in the body, whereas the **gonad** sex glands regulate sexual behavior and reproduction. The **adrenal glands** play a critical role in regulating the body's response to stress.

Two techniques used to study the brain involve either destroying a specific area of the brain (**lesioning**) or stimulating a specific brain area to see the effect. Researchers have developed several methods such as **CT**, **MEG**, **MRI**, **EEG**, **PET**, **fMRI**, or **TMS**.

The brain can be roughly divided into three sections: the forebrain, the midbrain, and the hindbrain. The many structures within each division are discussed, such as the **brainstem**, **medulla**, **reticular formation**, **limbic system**, **thalamus**, **hypothalamus**, **hippocampus**, **amygdala**, **cortex**, **cerebral hemispheres**, **corpus callosum**, **occipital lobes**, **parietal lobes**, **temporal lobes**, and **frontal lobes**. **Mirror neurons**, neurons that fire when we perform an action and also when we see someone else perform that action, may explain a great deal of the social learning that takes place in humans from infancy on. **Association areas** are the areas within each of the lobes that are responsible for "making sense" of all the incoming information. **Broca's area** is located in the left frontal lobe and **Wernicke's area** is located in the left temporal lobe; both play a role in language. The **cerebrum** is made up of the two cerebral hemispheres and the structures connecting them. **Split-brain research** helped scientists understand that the two cerebral hemispheres are not identical.

▲ Return to Chapter 2: Table of Contents

▼ CHANGES FROM THE FOURTH EDITION TO THE FIFTH EDITION

Chapter 2: The Biological Perspective

- Updated research on the importance of glial cells and myelin
- Moved *Psychology in the News* feature on misinterpretation of brain studies into the main narrative
- Added coverage of neurogenesis and expanded the discussion of neuroplasticity with research on new methods of repairing spinal cord damage and brain damage
- Enhanced discussion of epigenetics
- Incorporated new research on the relationship between sex hormones and cognitive changes during aging
- Added research on oxytocin's effects on the body and behavior
- Added a discussion of optogenetics, a developing area in brain stimulation
- Incorporated new research showing the connection between the cerebellum and disorders characterized by perceptual disturbances
- Included new research on the connection between the thalamus and task-specific regions of cortex
- Added new research suggesting activity in the amygdala impacts hippocampal neuroplasticity
- Revised the discussion of Broca's area and its connection to speech
- Added statistics on the rates of handedness
- New APA Goal 2 feature on Phineas Gage
- Incorporated new research in the "Applying Psychology to Everyday Life" feature on ADHD

▲ Return to Chapter 2: Table of Contents

▼ LECTURE LAUNCHERS AND DISCUSSION TOPICS

- 2.1 - Neurotransmitters: Chemical Communicators of the Nervous System
- 2.2 - Exceptions to the Rules
- 2.3 - The Glue of Life: Neuroglial Cells
- 2.4 - My Mother, Myself
- 2.5 - Brain Metaphors
- 2.6 - The Cranial Nerves
- 2.7 - Hormone Imbalances
- 2.8 - Psychophysiological Measurement
- 2.9 - Berger's Wave
- 2.10 - Lie Detectors 2.0
- 2.11 - Women, Men, and PETs
- 2.12 - Using fMRI and MEG to Study Phantom Limb Pain
- 2.13 - The Importance of a Wrinkled Cortex
- 2.14 - Brain's Bilingual Broca
- 2.15 - A New Look at Phineas Gage
- 2.16 - Freak Accidents and Brain Injuries
- 2.17 - Understanding Hemispheric Function
- 2.18 - Handedness, Eyedness, Footedness, Facedness
- 2.19 - Workplace Problems: Left-Handedness
- 2.20 - The Results of a Hemispherectomy

▲ Return to Chapter 2: Table of Contents

Lecture Launcher 2.1 – Neurotransmitters: Chemical Communicators of the Nervous System

In 1921, a scientist in Austria put two living hearts in a fluid bath that kept them beating. He stimulated the vagus nerve of one of the hearts. This is a bundle of neurons that serves the parasympathetic nervous system and causes a reduction in the heart's rate of beating. A substance was released by the nerve of the first heart and was transported through the fluid to the second heart. The second heart reduced its rate of beating. The substance released from the vagus nerve of the first heart was later identified as *acetylcholine*, one of the first neurotransmitters to be identified. Although many other neurotransmitters have now been identified, we continue to think of acetylcholine as one of the most important neurotransmitters. Curare is a poison that was discovered by South American Indians. They put it on the tips of the darts they shoot from their blowguns. Curare blocks acetylcholine receptors, and paralysis of internal organs results. The victim is unable to breathe, and eventually dies. A substance in the venom of black widow spiders stimulates release of acetylcholine at synapses. Botulism toxin, found in improperly canned foods, blocks release of acetylcholine at the synapses and has a deadly effect. It takes less than one millionth of a gram of this toxin to kill a person. A deficit of acetylcholine is associated with Alzheimer's disease, which afflicts a high percentage of older adults.

Many neurotransmitters have been identified in the years since 1921, and there is increasing evidence of their importance in human behavior. Psychoactive drugs affect consciousness because of their effects on synaptic transmission. For example, cocaine and the amphetamines prolong the action of certain neurotransmitters and opiates imitate the action of natural neuromodulators called the endorphins. It appears that the neurotransmitters dopamine, norepinephrine, and serotonin are associated with some of the most severe forms of mental illness.

There are probably only a few ounces of these substances in the body, but they may have a profound effect on mood, memory, perception, and behavior. Could intelligence be primarily a matter of having plenty of the right neurotransmitter at the right synapses?

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.2 - Exceptions to the Rules

In an introductory psychology class, students learn the basic rules that generally govern neuronal communication. In many cases, however, the exceptions to these rules may be as important as the rules themselves. Several of these exceptions are described below.

Rule #1: Neuron to neuron signaling is chemical, not electrical.

Exception: Gap junctions

Although it is generally the case that a neuron's electrical signal must first be converted to a chemical signal in order to excite or inhibit another neuron, this is not always the case. Some neurons have gap junctions, which connect their intracellular fluids. This means that the electrical signal can flow directly from one neuron to another. Unlike chemical synapses, most electrical synapses formed by gap junctions are bi-directional, meaning that electrical signals can travel in both directions through the gap junctions. Gap junctions also contain gates, which can be closed to prevent the electrical signal from being passed to the neighboring neuron.

Rule #2: Axons always synapse on dendrites.

Exception: Axo-axonic and axosomatic synapses

Axons can form synapses on all parts of a postsynaptic neuron. Synapses located on the soma (i.e., cell body) of a neuron are often inhibitory. In other words, transmitters released at these axosomatic synapses make it harder for the postsynaptic neuron to reach the threshold for generating an action potential. When an axon synapses on the axon of another neuron, it is called an axo-axonic synapse. Because these synapses usually occur near the end of the axon, they have no effect on whether the postsynaptic cell generates an action potential or not. Instead, axo-axonic synapses usually modulate how much neurotransmitter is released from the postsynaptic neuron.

Rule #3: Action potentials only travel in one direction.

Exception: Back-propagating action potentials

Action potentials begin at the axon hillock, where the axon emerges from the soma. From there, the action potential travels down the axon, and away from the soma. At the same time however, a back-propagating action potential can travel from the axon hillock, through the soma, and into the dendrites. Back-propagating action potentials are thought to affect the functioning of receptors located in the soma and dendrites.

Kandel, E., Schwartz, J., & Jessell, T. (2012). *Principles of neural science* (5th ed.). New York, NY: McGraw-Hill.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.3 - The Glue of Life: Neuroglial Cells

Glia is derived from the Greek word for glue and is an appropriate name for the cells that surround all

neurons, sealing them together. Glial cells outnumber neurons ten to one, and, although tiny in size, still make up half of the brain's bulk. Unlike neurons, glia do not possess excitable membranes and so cannot transmit information in the way neurons do. Yet so many thousands of cells must be there for some purpose.

Researchers studying the brain have suggested that glia can take up, manufacture, and release chemical transmitters, and so may help to maintain or regulate synaptic transmission. Other researchers suggest that glia can manufacture and possibly transmit other kinds of molecules, such as proteins. The anatomy of some glial cells is striking in this regard, for they seem to form a conduit between blood vessels and neurons, and so may bring nourishment to the neurons. It is thought that these cells may have important functions during prenatal development and recovery from brain injury. One role of the glia is known definitely: certain kinds of glia, called by the tongue-twisting name of oligodendroglia, form the myelin sheath that insulates central nervous system axons and speeds conduction of the nerve impulse. A counterpart called a Schwann cell performs the same role for the neurons that make up peripheral nerves.

The study of glia is difficult because these tiny cells are inextricably entwined with neurons. As the most numerous type of cell in the brain, their potential importance is vast, and investigation of their function seems likely to yield exciting results in the near future.

[**► Return to Lecture Guide**](#)

[**◄ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2**](#)

[**▲ Return to Chapter 2: Table of Contents**](#)

Lecture Launcher 2.4 - My Mother, Myself

Many people feel that their mothers are “a part of them.” Recent findings suggest that there may be considerable truth to that phrase.

Many adults apparently still have cells in their bodies that they picked up from their mothers during the gestation period. Similarly, many mothers still have cells in their bodies that came from their own children during pregnancy. Technically speaking, these “guest cells” are actually the product of stem cells that got planted in the “host’s” body and started reproducing decades later. And, technically speaking, there aren’t too many of them. Some estimates put the number of foreign cells at less than one in a million, a comforting thought for anyone conjuring up images of parasitic offspring or alien-like entities living happily rent-free.

The meaning of these *microchimeras* is less clear. There is some evidence that these cells might contribute to autoimmune diseases, although there is also speculation that these cells might confer a health benefit. Because this area of study is relatively young, there remain more questions than answers (such as, what about women who have cells from both their mothers and their own offspring?). It’s comforting to know, though, that in some small way a parent is always with us.

Boddy, A. M., Fortunato, A., Sayres, M. W., & Aktipis, A. (2015). Fetal microchimerism and maternal health: A review and evolutionary analysis of cooperation and conflict beyond the womb. *Bioessays*, 37(10), 1106-1118.

Gammill, H. S., & Nelson, J. L. (2010). Naturally acquired microchimerism. *International Journal of Developmental Biology*, 54, 531-543.

Ritter, M. (2003, May 9). Moms, you’ve got a little of your kids inside you. *Austin American-Statesman*, A1, A16.

Shute, N. (2010, Apr. 3). Beyond birth: A child’s cells may help or harm a mother long after delivery. *Scientific American*.
<http://www.microchimerism.org/>

[**► Return to Lecture Guide**](#)

[**◄ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2**](#)

[**▲ Return to Chapter 2: Table of Contents**](#)

Lecture Launcher 2.5 - Brain Metaphors

Metaphors are a toolbox of handiness in psychology, because they help us to understand systems that aren't directly observable through reference to things that are more familiar and perhaps better understood (Weiner, 1991). Our understanding of the human brain and its activity has been helped through a reliance on metaphor. The metaphors used, however, have changed over time.

- *Hydraulic models.* Thinkers such as Galen and Descartes described the brain as a pneumatic/hydraulic system, relying on the “new-fangled” plumbing systems dominant during their lifetimes. Galen, for example, believed that the liver generated “spirits” or gases that flowed to the brain, where they then formed “animal spirits” that flowed throughout the nervous system. Descartes expanded on this view, adding that the pineal gland (the supposed seat of the soul) acted on the animal spirits to direct reasoning and other behaviors. In short, the brain was a septic tank; storing, mixing, and directing the flow of spirit gases throughout the body for the purposes of behavior and action.

- *Mechanical and telephone models.* With the advent of new technology came new metaphors for the brain. During the Industrial Revolution machine metaphors dominated, and in particular the brain was conceived as a complex mechanical apparatus involving (metaphorical) levers, gears, trip hammers, and pulleys. During the 1920s, the brain developed into a slightly more sophisticated machine resembling a switchboard; the new technology of the telephone provided a new metaphor. Inputs, patch cords, outputs, and busy signals (though no “call waiting”) dominated explanations of brain activity. This metaphor, however, faltered by viewing the brain as a system that shut down periodically, as when no one was dialing a number. We now know, of course, that the brain is continually active.

- *Computer models.* Starting in the late 1950's, metaphors for the brain relied on computer technology. Input, output, memory, storage, information processing, and circuitry were all terms that seemed equally suited to talking about computer chips or neurons. Although perhaps a better metaphor than plumbing or telephones, the computer model eventually showed its shortcomings. As a descriptive device, however, this metaphor can at least suggest limits in our understanding and point the way to profitable areas of research.

Weiner, B. (1991). Metaphors in motivation and attribution. *American Psychologist*, 46, 921-930.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.6 - The Cranial Nerves

The textbook discusses various divisions of the nervous system. You may want to add a description of the cranial nerves to your outline of the nervous system. Although the function of the cranial nerves is not different from that of the sensory and motor nerves in the spinal cord, they do not enter and leave the brain through the spinal cord. There are twelve cranial nerves (numbered 1 to 12 and ordered from the front to the back of the brain) that primarily transmit sensory information and control motor movements of the face and head. The twelve cranial nerves are:

1. *Olfactory.* A sensory nerve that transmits odor information from the olfactory receptors to the brain.
2. *Optic.* A sensory nerve that transmits information from the retina to the brain.
3. *Oculomotor.* A motor nerve that controls eye movements, the iris (and therefore pupil size), lens accommodation, and tear production.

4. *Trochlear*. A motor nerve that is also involved in controlling eye movements.
5. *Trigeminal*. A sensory and motor nerve that conveys somatosensory information from receptors in the face and head and controls muscles involved in chewing.
6. *Abducens*. Another motor nerve involved in controlling eye movements.
7. *Facial*. Conveys sensory information and controls motor and parasympathetic functions associated with facial muscles, taste, and the salivary glands.
8. *Auditory-vestibular*. A sensory nerve with two branches, one of which transmits information from the auditory receptors in the cochlea and the other conveys information concerning balance from the vestibular receptors in the inner ear.
9. *Glossopharyngeal*. This nerve conveys sensory information and controls motor and parasympathetic functions associated with the taste receptors, throat muscles, and salivary glands.
10. *Vagus*. Primarily transmits sensory information and controls autonomic functions of the internal organs in the thoracic and abdominal cavities.
11. *Spinal accessory*. A motor nerve that controls head and neck muscles.
12. *Hypoglossal*. A motor nerve that controls tongue and neck muscles.

As is their custom, medical students have developed several mnemonics for memorizing the cranial nerves. Some of the family-friendly ones include:

**On Old Olympus' Tiny Tops, A Friendly Viking Grew Vines And Hops
 Oh Once One Takes The Anatomy Final Very Good Vacations Are Heavenly
 One Of Our Two Timing Adults Found Very Good Values At Home
 On Occasion Our Trusty Truck Acts Funny. Very Good Vehicle Any How
 Orlando's Overweight Octopuses Try To Avoid Fuddrucker's And Grabbing Vienna Sausage Hamburgers
 On Our Overseas Trip To Argentina Found Very Grand Villas And Huts**

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.7 - Hormone Imbalances

Various problems are caused by imbalances within the endocrine system. The following disorders and medical problems are associated with abnormal levels within the pituitary, thyroid, and adrenal glands.

Pituitary malfunctions

Hypopituitary Dwarfism

If the pituitary secretes too little of its growth hormone during childhood, the person will be very small, although normally proportioned.

Giantism

If the pituitary gland over-secretes the growth hormone while a child is still in the growth period, the long bones of the body in the legs and other areas grow very, very long—a height of 9 feet is not unheard of. The organs of the body also increase in size, and the person may have health problems associated with both the extreme height and the organ size.

Acromegaly

If the over-secretion of the growth hormone happens after the major growth period is ended, the person's long bones will not get longer, but the bones in the face, hands, and feet will increase in size, producing abnormally large hands, feet, and facial bone structure. The famous wrestler/actor, Andre the Giant (Andre Rousimoff), had this condition, as did the great actor Rondo Hatton.

Thyroid malfunctions

Hypothyroidism

In hypothyroidism, the thyroid does not secrete enough thyroxin, resulting in a slower than normal metabolism. The person with this condition will feel sluggish and lethargic, have little energy, and tend to be obese.

Hyperthyroidism

In hyperthyroidism, the thyroid secretes too much thyroxin, resulting in an overly active metabolism. This person will be thin, nervous, tense, and excitable. He or she will also be able to eat large quantities of food without gaining weight (oh, if only we came equipped with thyroid control knobs!).

Adrenal Gland Malfunctions

Among the disorders that can result from malfunctioning of the adrenal glands are Addison's Disease (which is caused by adrenal insufficiency) and Cushing's Syndrome (caused by elevated levels of cortisol). In the former, fatigue, low blood pressure, weight loss, nausea, diarrhea, and muscle weakness are some of the symptoms, whereas for the latter, obesity, high blood pressure, a "moon" face, and poor healing of skin wounds is common. John F. Kennedy, Helen Reddy, and (perhaps) Osama bin Laden were well-known Addisonians.

If there is a problem with over-secretion of the sex hormones in the adrenals, virilism and premature puberty are possible problems. Virilism results in women with beards on their faces and men with exceptionally low, deep voices. Premature puberty, or full sexual development while still a child, is a result of too many sex hormones during childhood. (Puberty is considered premature if it occurs before the age of 8 in girls and 9 in boys.) Treatment is possible using hormones to control the appearance of symptoms, but must begin early in the disorder.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.8 - Psychophysiological Measurement

There are various strategies for measuring activity in the brain, especially recently developed techniques such as PET, TMS, or MRI. There are, of course, other bodily systems and other techniques for measuring them, many of which rely on the electrophysical activity of the body.

- *EMG—Electromyography.* An electromyogram records the action potential given off by contracting muscle fibers. A common example is the recording of facial EMG, in which either inserted electrodes or surface electrodes record the activity of muscles as they pose various expressions.

- *EGG—Electrogastrography.* Electrogastrograms provide a record of smooth muscle activity in the abdomen. The contractions of the stomach or intestines, for example, can be measured by comparing the readings from a surface electrode attached to the abdomen with those of an electrode attached to the forearm. In the special case of measuring contractions in the esophagus, surface electrodes are attached to a balloon, which is "swallowed" by the person being measured. EGG may be used successfully to gain information about fear, anxiety, or other emotional states.

- *EOG—Electrooculography*. Readings from electrodes placed around the posterior of the eyes are the basis for EOG. Electrical signals result from small saccadic eye movements as well as more gross movements that can be directly observed. EOG can be used for measuring rapid eye movements during sleep.

- *EKG—Electrocardiography*. EKG records changes in electrical potential associated with the heartbeat. Electrodes are placed at various locations on the body, and their recordings yield five waves that can be analyzed: P-waves, Q-waves, R-waves, S-waves, and T-waves. EKG may be used by psychologists to supplement observations relevant to stress, heart disease, or Type A behavior patterns.

- *EDA—Electrodermal Activity*. Formerly called *galvanic skin response*, *skin resistance*, and *skin conductance*, EDA refers to the electrical activity of the skin. As activity in the sympathetic nervous system increases it causes the eccrine glands to produce sweat. This activity of the eccrine glands can be measured by EDA, regardless of whether or not sweat actually rises to the skin surface. The folklore of “sweaty palms” associated with a liar might be measured using this technique.

- *EEG—Electroencephalography*. As discussed in the text, EEG provides information about the electrical activity of the brain, as recorded by surface electrodes attached to the scalp. EEG has been used in a variety of ways to gather information about brain activity under a wide range of circumstances.

- *Pneumography*. Pneumographs measure the frequency and amplitude of breathing, and are obtained through a relatively straightforward procedure. A rubber tube placed around the chest expands and contracts in response to the person’s inhalations and exhalations. These changes can then be recorded with either an ink pen or electrical signal.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.9 - Berger’s Wave

Ask if anyone knows what is meant by the term *Berger’s wave*. Explain that the study of electrical activity in the brain was once limited to studies in which different kinds of measuring devices were attached to the exposed brains of animals. Studies involving humans were rare; researchers could only measure the electrical activity of the living human brain in individuals who had genetic defects of their skull bones that caused the skin of their scalps to be in direct contact with the surfaces of their brains. Yuck!

All this changed when a German physicist named Hans Berger, after several years of painstaking research, discovered that it was possible to amplify and measure the electrical activity of the brain by attaching special electrodes to the scalp which, in turn, sent impulses to a machine that graphed them. In his research, Berger discovered several types of waves, one of which he called the “alpha” wave for no other reason than being the first one he discovered (“alpha” is the first letter of the Greek alphabet). He kept his research a secret until he published an article about it in 1929. The alpha wave is also sometimes called *Berger’s wave* in honor of Berger’s discovery.

Obviously, Berger achieved one of the most important discoveries in the history of neuroscience. However, his life was not a happy one. Shortly after his article was published, the Nazis rose to power in Germany, which greatly distressed him. In addition, his work wasn’t valued in Germany; he was far better known in the United States. As a result, Berger fell into a deep depression in 1941 and hanged himself.

Gloor, P. (1969). Hans Berger and the discovery of the electroencephalogram. *Electroencephalography and Clinical Neurophysiology*: Supplement 28, 1–36.

Millett, D. (2001). Hans Berger: From psychic energy to the EEG. *Perspectives in Biological Medicine*, 44 (4), 522–542

Wiedemann, H. R. (1994). Hans Berger (1873-1941). *European Journal of Pediatrics*, 153(10), 705.

► [Return to Lecture Guide](#)

[◀ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

[▲ Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.10 - Lie Detectors 2.0

A staple of police and lawyer television shows is the “lie detector” scene, in which the suspect is hooked up to polygraph machine and asked a series of questions about a crime. As the questions are asked, the needles on the polygraph record the suspect’s heart rate, breathing, skin conductance, and other physiological responses to the questions. Polygraph machines have been used in this way by various law enforcement agencies for many years. The principle behind the test is that the act of lying causes an involuntary change in the autonomic nervous system, which can be detected by the polygraph. The accuracy of polygraph machines, however, is controversial, and in many courts they are inadmissible evidence. More recently, some researchers have tried to create a new generation of lie detectors, which can measure activity in the brain directly. These techniques look for patterns in the brain that, at least in theory, correlate with lying.

One technique that might be adapted to lie-detection is electroencephalography, more commonly referred to as EEG. During an EEG recording, electrodes are placed at various locations on the scalp. These electrodes are capable of picking up the electrical activity produced by neurons located in different parts of the brain. While the activity of individual neurons cannot be identified, the patterns of electrical activity produced by thousands of neurons working together can be a sign that the brain is functioning in a particular way. One way EEGs may be useful as lie detectors, is by identifying event-related potentials or (ERPs). An ERP is a brief electrical change that occurs at a reliable time point relative to a specific event. For example, it has been found that 300 to 500 ms after a person has been shown something that is unexpected or novel, there is a brief electrical change in that person’s EEG. Theoretically, this ERP could be used to determine if a subject has previous knowledge of a piece of evidence. For instance, an ERP occurring 300 ms after being shown a picture of the murder weapon might indicate that the suspect had not seen the murder weapon before.

More recently, fMRIs have been suggested as potential lie-detectors. fMRI, or functional magnetic resonance imaging, works by detecting the increase in blood flow to more active regions in the brain. This is not to be confused with structural MRIs, which can only create an image of tissues, bones, etc. When a person performs a task in an fMRI, adding two numbers together for example, the brain regions required to perform the task will become active. This activity will cause a change in blood flow, which the fMRI can detect. It is possible that, because different brain regions are involved in recounting an actual event than are involved in making up a story, an fMRI is capable of determining whether someone is lying or telling the truth. Some researchers have found that, even if a lie is well rehearsed, it still appears to activate different brain regions than telling the truth does.

Despite media interest in new forms of lie-detection, many experts agree that the EEG and fMRI approaches currently suffer from the same issues that polygraphs do. For example, although the newer techniques measure brain activity much more directly, there is concern about their reliability. While certain brain activity might suggest that a person is lying, unless the technology can be made almost 100% accurate, innocent people may be accused of crimes they did not commit. Also, it is unclear whether people could find ways to “trick” the machines by performing certain mental tasks during testing. Until these questions can be answered, it is unlikely that the polygraph will be replaced anytime soon.

Farah, M. J., Hutchinson, J. B., Phelps, E. A., & Wagner, A. D. (2014). Functional MRI-based lie detection: Scientific and societal challenges. *Nature Reviews | Neuroscience*, 15, 123-131.

Ganis, G., Kosslyn, S., Stose, S., Thompson, W., & Yurgelun-Todd, D. (2003). Neural correlates of different types of deception: An fMRI investigation. *Cerebral Cortex*, 13(8), 830-836.

Wolpe, P., Foster K., & Langleben D. (2005). Emerging neurotechnologies for lie-detection: Promises and perils. *The American Journal of Bioethics*, 5(2), 39-49.

[► Return to Lecture Guide](#)

[◀ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

[▲ Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.11 - Women, Men, and PETs

The 1990s were dubbed “the decade of the brain,” and it is true that remarkable advances have been made by the neurosciences in discovering how the brain operates. Several studies suggest that the operation of men’s and women’s brains may differ in significant ways.

For example, Ruben Gur and his colleagues at the University of Pennsylvania recorded PET scans of men and women who were asked to think of nothing in particular. That is, the research participants were instructed to relax and let their brains idle as they exerted as little mental effort as possible. The researchers found that for most participants the task was difficult to complete; PET scans revealed that these idle minds nonetheless hummed with activity. The locus of that activity, however, differed across the sexes. Men’s brains often showed activity in the limbic system, whereas women often showed activity in the posterior cingulate gyrus. The meaning of these differences is difficult to interpret; the difficulty is compounded by the 13 men and 4 women who showed patterns of activity characteristic of their opposite sex peers. As an early peek into the brain, however, they hint that the centers of activity for “blank” brains differ for women and men.

In a separate study, researchers at the University of California, Irvine, asked 22 men and 22 women to solve SAT math problems while undergoing a PET scan. Half of each group had SAT math scores above 700, whereas the other half had scores below 540. The temporal lobes of the 700+ men showed heightened activity during the math task, although this was not true for the women; the 700+ women’s temporal lobes were no more intensively used than those of the 540-group women. Richard Haier, who helped lead the study, speculates that women in the top group might be using their brains more efficiently than women in the average-scoring group. More generally, although both men and women did well at the task, their brains were operating differently to accomplish it.

Ruben and Raquel Gur also studied men’s and women’s brains in response to emotional expressions (Erwin et al., 1992). Shown pictures of either happy or sad faces, both men and women were quite adept at spotting happiness. Women, however, could identify sadness about 90% of the time, regardless of whether it was on the face of a man or a woman. By comparison, men were accurate in spotting sadness 90% of the time on a man’s face, but only 70% of the time if the expression was posed by a woman. Once again, PET scans revealed that women’s brains didn’t have to work as hard at this task as did men’s; in fact, women’s limbic systems were less active than the limbic systems of the poor-scoring men.

There are a number of other differences between women’s and men’s brains. Women tend to have a larger corpus callosum than men, for example. Women may have a higher concentration of neurons in their cortexes than men, for another. But the meaning behind these differences is a matter far from decided.

Begley, S. (1995, March 27). Gray matters. *Newsweek*, pp. 48-54.

Erwin, R. J., Gur, R. C., Gur, R. E., Skolnick, B. et al. (1992). Facial emotion discrimination I: Task construction and behavioural findings in normal subjects. *Psychiatry Research*, 42, 231-240.

► Return to Lecture Guide

◄ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2

▲ Return to Chapter 2: Table of Contents

Lecture Launcher 2.12 – Using fMRI and MEG to Study Phantom Limb Pain

The concept of *pain sensation* means different things to different people. Many students are aware of phantom pain sensations and are actually very curious as to what it is. Medical professionals have recorded many cases of what has come to be called “phantom limbs.” Phantom limb phenomenon occurs

when a person who has had an amputation of some body part, such as an arm or leg, reports “feeling” sensations from the now-missing limb. Phantom limb refers to the subjective sensory awareness of an amputated body part, and may include numbness, itchiness, temperature, posture, volume, or movement. For example, one man whose left arm was amputated just above the elbow during a horrific car accident claimed that he could still feel the arm as a kind of ghostly presence. He could feel himself wiggling non-existent fingers and “grabbing” objects that would have been in his reach had his arm still been there (Ramachandran & Blakeslee, 1998). Phantom sensations may take years to fade, and usually do so from the end of the limb up to the body—in other words, one’s phantom arm seems to get shorter and shorter until it can no longer be felt. In addition to legs and arms there have been cases of phantom breasts, bladders, rectums, vision, hearing, and internal organs.

Phantom limb pain refers to the specific case of painful sensations that appear to reside in the amputated body part. Patients have variously reported pins-and-needles sensations, burning sensations, shooting pains that seem to travel up and down the limb, or cramps, as though the severed limb was in an uncomfortable and unnatural position. Many amputees often experience several types of pain; others report that the sensations are unlike other pain they’ve experienced. Unfortunately, some estimates suggest that over 70 percent of amputees still experience intense pain, even 25 years after amputation. Most treatments for phantom limb pain (there are over 50 types of therapy) help only about 7 percent of sufferers.

What causes these phantom sensations? A recent study has shed light on the causes of phantom limb sensations. Researchers at Humboldt University in Berlin suggest that the most severe type of this pain occurs in amputees whose brains undergo extensive sensory reorganization. Magnetic responses were measured in the brains of 13 arm amputees in response to light pressure on their intact thumbs, pinkies, lower lips, and chins. These responses were then mapped onto the somatosensory cortex controlling that side of the body. Because of the brain’s contralateral control over the body, the researchers were able to estimate the location of the somatosensory sites for the missing limb. They found that those amputees who reported the most phantom limb pain also showed the greatest cortical reorganization. Somatosensory areas for the face encroached into regions previously reserved for the amputated fingers.

Renowned neuroscientist Dr. V. S. Ramachandran has investigated many cases of phantom limb sensations in his career. He believes that examination of people who experience these phenomena, using the non-invasive techniques of magnetoencephalograms and functional MRIs, can teach us much about the relationship between sensory experience and consciousness. Researchers have long known that touching certain points on the stump of the amputation (and in some cases on the person’s face) can produce phantom sensations in a missing arm or fingers (Ramachandran & Hirstein, 1998). Older explanations of phantom limb sensations have called it an illusion brought on by the irritation of the nerve endings in the stump due to scar tissue. But using anesthesia on the stump does not remove the phantom limb sensations or the pain experienced by some patients in the missing limb, so that explanation is not adequate. Ramachandran and colleagues suggest instead that phantom limb sensations may occur because areas of the face and body near the stump “take over” the nerve functions that were once in the control of the living limb, creating the false impression that the limb is still there, feeling and moving. This “remapping” of the limb functions, together with the sensations from the neurons ending at the stump and the person’s mental “body image” work together to produce phantom limb sensations.

Although these findings do not by themselves solve the riddle of phantom limb pain, they do offer avenues for future research. For example, damage to the nervous system may cause a strengthening of connections between somatosensory cells and the formation of new ones. Phantom limb pain may result due to an imbalance of pain messages from other parts of the brain. As another possibility, pain may result from a remapping of somatosensory areas that infringes on pain centers close by.

Boas, R. A., Schug, S. A., & Acland, R. H. (1993). Perineal pain after rectal amputation: A 5-year follow-up. *Pain*, 52, 67–70.
Bower, B. (1995). Brain changes linked to phantom-limb pain. *Science News*, 147, 357.
Brena, S. F., & Sammons, E. E. (1979). Phantom urinary bladder pain – Case report. *Pain*, 7, 197–201.

Bressler, B., Cohen, S. I., & Magnussen, F. (1955). Bilateral breast phantom and breast phantom pain. *Journal of Nervous and Mental Disease*, 122, 315–320.

Dorpat, T. L. (1971). Phantom sensations of internal organs. *Comprehensive Psychiatry*, 12, 27–35.

Katz, J. (1993). The reality of phantom limbs. *Motivation and Emotion*, 17, 147–179.

Ramachandran, V. S., & Blakeslee, S. (1998). *Phantoms in the Brain*. William Morrow, N.Y.

Ramachandran, V. S., & Hirstein, W. (1998). The perception of phantom limbs: The D. O. Hebb lecture. *Brain*, 121, 1603–1630.

Shreeve, J. (1993, June). Touching the phantom. *Discover*, 35–42.

[► Return to Lecture Guide](#)

[◄ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

[▲ Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.13 - The Importance of a Wrinkled Cortex

At the beginning of your lecture on the structure and function of the brain, ask students to explain why the cerebral cortex is wrinkled. There are always a few students who correctly answer that the wrinkled appearance of the cerebral cortex allows it to have a greater surface area while fitting in a relatively small space (i.e., the head). To demonstrate this point to your class, hold a plain, white sheet of paper in your hand and then crumple it into a small, wrinkled ball. Note that the paper retains the same surface area, yet is now much smaller and is able to fit into a much smaller space, such as your hand. You can then mention that the brain's actual surface area, if flattened out, would be roughly the size of a newspaper page. Laughs usually erupt when the class imagines what our heads would look like if we had to accommodate an unwrinkled, newspaper-sized cerebral cortex!

[► Return to Lecture Guide](#)

[◄ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

[▲ Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.14 - Brain's Bilingual Broca

Se potete parlare Italiano, allora potete capire questa sentenza. Of course, if you only speak English, you probably only understand *this* sentence. If you speak both languages, then by this point in the paragraph you should be really bored.

Bilingual speakers who come to their bilingualism in different ways show different patterns of brain activity. Joy Hirsch of Memorial Sloan-Kettering Cancer Center in New York and her colleagues monitored the activity in Broca's area in the brains of bilingual speakers who acquired their second language starting in infancy, and compared it to the activity of bilingual speakers who adopted a second language in their teens. Participants were asked to silently recite brief descriptions of an event from the previous day, first in one language and then in the other. A functional magnetic resonance image (fMRI) was taken during this task. All of the 12 adult speakers were equally fluent in both languages, used both languages equally often, and represented speakers of English, French, and Turkish, among other tongues.

Hirsch and her colleagues found that among the infancy-trained speakers, the same region of Broca's area was active, regardless of the language they used. Among the teenage-trained speakers, however, a different region of Broca's area was activated when using the acquired language. Similar results were found in Wernicke's area in both groups. Although the full meaning of these results is a matter of some debate (do they reflect sensitivity in Broca's area to language exposure, or pronounced differences in adult versus childhood language learning?), they nonetheless reveal an intriguing link between *la testa e le parole*.

Bower, B. (1997, July 12). Brains show signs of two bilingual roads. *Science News*, 152, 23.

[► Return to Lecture Guide](#)

[◄ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

Lecture Launcher 2.15 - A New Look at Phineas Gage

For over 30 years, Jack and Beverly Wilgus had a daguerreotype portrait -- a type of early photograph -- of a well dressed young man with one eye closed. Because the photo showed the young man holding what appeared to be part of a harpoon, the Wilguses believed that the man was a 19th century whaler who had lost his eye, perhaps in a whaling accident. It was only after a copy of the portrait was posted online that the couple was told that the object in the man's hands did not appear to be a harpoon. Then, in 2008, a person viewing the image online posted a comment that the young man may be Phineas Gage, making the "harpoon" the infamous tamping rod that was blasted through his skull and brain. By carefully examining the rod in the daguerreotype, and by comparing the young man's face to the cast made of Gage's head after his death, the Wilguses were able to confirm that the portrait is almost certainly that of Phineas Gage, made sometime after his accident. Importantly, this is the only known photograph of the man who became one of the most famous case studies in psychology.

One of the consequences of the portrait's discovery has been a renewed debate about how Gage's injuries affected his personality and behavior. Many psychology textbooks explain that the accident left Gage a permanently changed man following the accident, with his once well-balanced, gregarious, and hard-working personality replaced with profane, inconsiderate, and impulsive behavior for the rest of his life. This, however, is not necessarily supported by the few original sources researchers have to go on. For example, although the evidence clearly indicates that Gage had major psychological changes for a period after his accident, we also know that Gage later spent many years driving stagecoaches before he died in 1860, 12 years after the accident. Many have questioned whether the post-accident Phineas Gage commonly described in introductory psychology classes could have performed the tasks required to drive a stagecoach, interact with passengers, and be reliable enough to maintain employment for long periods at a time. Does this indicate that many of the psychological changes Gage suffered were temporary? Certainly the newly discovered daguerreotype of a healthy looking and well-kept Phineas Gage lends further support to the idea that Phineas was able to largely recover from his accident, both physically and mentally. If true, this may mean that the case of Phineas Gage may be as much a story about the incredible plasticity of the brain and its ability to compensate for the loss of specific brain regions, as it is about the localization of specific functions.

The newly discovered portrait of Phineas Gage can be found at: brightbytes.com/phineasgage or by searching the Internet for "Phineas Gage daguerreotype."

Jarrett, C. & Sutton, J. (2009, July 20). Face to face with Phineas Gage. Retrieved December 9, 2009, from, <http://www.thepsychologist.org.uk/blog/11/blogpost.cfm?threadid=1035&catid=48>
Macmillan, M. (2008). Phineas Gage—Unravelling the myth. *The Psychologist*, 21(9), 828-831.

► Return to Lecture Guide

◄ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2

▲ Return to Chapter 2: Table of Contents

Lecture Launcher 2.16 - Freak Accidents and Brain Injuries

Students may be interested in the unusual cases of individuals who experience bizarre brain injuries due to freak accidents with nail guns. The most fascinating example involved Isidro Mejias, a construction worker in Southern California, who had six nails driven into his head when he fell from a roof onto his coworker who was using a nail gun. (X-ray images of the embedded nails can be found at several sites on the Internet.) Incredibly, none of the nails caused serious damage to Mejia's brain. One nail lodged near his spinal cord, and another came very close to his brain stem. Immediate surgery and treatment

with antibiotics prevented deadly infections that could have been caused by the nails. In a similar accident, a construction worker in Colorado ended up with a nail lodged in his head due to a nail gun mishap. Unlike Mejia, Patrick Lawler didn't realize he had a nail in his head for six days. The nail was discovered when he visited a dentist due to a "toothache." It appears that Lawler fired a nail into the roof of his mouth. The nail barely missed his brain and the back of his eye.

Nail Gun /Victim Lives. *Current Science*, A Weekly Reader publication, Sept. 10, 2004, v90 (1), Page 14.
<http://www.msnbc.msn.com/id/4909827/>
<http://www.summitdaily.com/article/20050119/NEWS/50119002/0/FRONTPAGE>

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.17 - Understanding Hemispheric Function

A variation on the rather dubious statement that "we only use one-tenth of our brain" is that "we only use one-half (hemisphere) of our brain." Research suggests that each cerebral hemisphere is specialized to perform certain tasks (e.g., left hemisphere/language; right hemisphere/ visuospatial relationships), with the abilities of one hemisphere complementary being to the other. From this came numerous distortions, oversimplifications, and unwarranted extensions, many of which are discussed in two interesting reviews of this trend toward "dichomania" (Corballis, 1980; Levy, 1985). For example, the left hemisphere has been described variously as logical, intellectual, deductive, convergent, and "Western," while the right hemisphere has been described as intuitive or creative, sensuous, imaginative, divergent, and "Eastern." Even complex tasks are described as right- or left-hemispheric because of their language component. In every individual one hemisphere supposedly dominates, affecting that person's mode of thought, skills, and approach to life. One commonly cited, but questionable test for dominance is to note the direction of gaze when a person is asked a question (left gaze signaling right hemisphere activity; right gaze showing left hemisphere activity). Advertisements have claimed that artistic abilities can be improved if the right hemisphere is freed, and the public schools have been blamed for stifling creativity by emphasizing left-hemisphere skills and by neglecting to teach the children's right hemisphere.

Corballis and Levy explode these myths and trace their development. In reality, the two hemispheres are quite similar and can function remarkably well even if separated by split-brain surgery. Each hemisphere does have specialized abilities, but the two hemispheres work together in all complex tasks. For example, writing a story involves left-hemispheric input concerning syntax, but right-hemispheric input for developing an integrated structure and for using humor or metaphor. The left hemisphere is not the sole determinant of logic, nor is the right hemisphere essential for creativity. Disturbances of logic are more prevalent with right-hemisphere damage, and creativity is not necessarily affected. Although one hemisphere can be somewhat more active than the other, no individual is purely "right brained" or "left brained." Also, eye movement and hemispheric activity patterns poorly correlate with cognitive style or occupation. Finally, because of the coordinated, interactive manner of functioning of both hemispheres, educating or using only the right or left hemisphere is impossible (without split-brain surgery).

Corballis, M.C. (1980). Laterality and myth. *American Psychologist*, 35, 284–295.
Levy, J. (1985). Right brain, left brain: Fact or fiction? *Psychology Today*, 19, 38–45.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.18 – Handedness, Eyedness, Footedness, Facedness

Although the title sounds like a Dr. Seuss rhyme, it actually denotes something sensible to neuropsychologists. Most people are familiar with the concept of handedness. The human population is

Copyright © 2017 Pearson Education. All rights reserved.

distributed across many people who are adept at using their right hands for most tasks, some who have greater skill using the left hand, and a smaller proportion of those who are equally skilled using either hand (or who alternate hands for certain tasks). The concepts of footedness, leggedness, eyedness, and facedness may be less familiar to the layperson, although they stem from the same principle as handedness.

The basis of these distinctions lies in the concept of laterality. Just as the cerebral hemispheres show specialization (e.g., left hemisphere language functions, right hemisphere visual-spatial functions), so too are there preferences or asymmetries in other body regions. The concept of eyedness, then, refers to the preference for using one eye over another, such as when squinting to site down the crosshairs of a rifle or to thread a needle. Footedness and leggedness similarly refer to a preference for one limb over the other; drummers and soccer players will attest to the importance of being equally adept at using either foot, and to the difficulty in achieving that. Finally, facedness refers to the strength with which information is conveyed by the right or left side of the face. It has been suggested that verbal information shows a right-face bias whereas emotional expressions are more strongly shown on the left side of the face, although these conclusions remain somewhat controversial.

Why are these distinctions useful? They play their largest role in the areas of sensation and perception, engineering psychology, and neuropsychology. Studies of reaction time, human-machine interaction, ergonomic design, and so on, take into account the preferences and dominance of some body systems over others. In the case of facedness and emotional expression, researchers are working to illuminate the link between facial expressions and cerebral laterality. Given the right hemisphere's greater role in emotional activities, the contralateral control between the right hemisphere and the left hemiface becomes an important proving ground for investigating both brain functions and the qualities of expression.

Borod, J. C., Caron, H. S., & Koff, E. (1981). Asymmetry of facial expression related to handedness, footedness, and eyedness: A quantitative study. *Cortex*, 17, 381-390.

Ekman, P., Hagar, C. J., & Friesen, W. V. (1981). The symmetry of emotional and deliberate facial actions. *Psychophysiology*, 18, 101-106.

Friedlander, W. J. (1971). Some aspects of eyedness. *Cortex*, 7, 357-371.

McGuigan, F. J. (1994). *Biological psychology: A cybernetic science*. Englewood Cliffs, NJ: Prentice Hall.

Sackheim, H. A., Gur, R. C., & Saucy, M. C. (1978). Emotions are expressed more intensely on the left side of the face. *Science*, 202, 434-436.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Lecture Launcher 2.19 - Workplace Problems: Left-Handedness

Within Canada and the United States, there are approximately 33 million people who are left handed. This presents a severe detriment to the workplace. It has been shown that left handed individuals are more likely to have accidents at work than are right handed individuals, in fact 25% more likely (and if they are working with tools and machinery, 51% more likely). Accommodations such as being able to rearrange the work area and having tools available that are either left- or right-hand adapted would make the workplace a safer place to be. Have students suggest ways that the workplace could be made safer or even what could be done in the classroom that would make it easier for students who are left handed to take notes or tests. What about the mouse on computers? The mouse is actually made for people who are right handed. How adaptable must a left handed person become in order not to be frustrated by using a right handed mouse?

Gunsch, D. For Your Information: Left-handed workers struggle in a right-handed work world. *Personnel Journal*, 93, 23-24.
<http://www.cnn.com/2015/11/03/health/being-left-handed-health-impact/>

► [Return to Lecture Guide](#)

◄ [Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2](#)

Lecture Launcher 2.20 - The Results of a Hemispherectomy

When Matthew was 6 years old, surgeons removed half of his brain.

His first three years of life were completely normal. Just before he turned 4, however, Matthew began to experience seizures, which did not respond to drug treatment. The seizures were both life threatening and frequent (as often as every three minutes). The eventual diagnosis was Rasmussen's encephalitis, a rare and incurable condition of unknown origin.

The surgery, a hemispherectomy, was performed at Johns Hopkins Hospital in Baltimore. A few dozen such operations are performed each year in the United States, usually as a treatment for Rasmussen's and for forms of epilepsy that destroy the cortex but do not cross the corpus callosum. After surgeons removed Matthew's left hemisphere, the empty space quickly filled with cerebrospinal fluid.

The surgery left a scar that runs along one ear and disappears under his hair; however, his face has no lopsidedness. The only other visible effects of the operation are a slight limp and limited use of his right arm and hand. Matthew has no right peripheral vision in either eye. He undergoes weekly speech and language therapy sessions. For example, a therapist displays cards that might say "fast things" and Matt must name as many fast things as he can in 20 seconds. He does not offer as many examples as other children his age. However, he is making progress in the use of language, perhaps as a result of fostering and accelerating the growth of dendrites.

The case of Matthew indicates the brain's remarkable plasticity. Furthermore, it is interesting to note that Matt's personality never changed through the seizures and surgery.

Boyle, M. (1997, August 1). Surgery to remove half of brain reduces seizures. *Austin American-Statesman*, A18.

Rasmussen T, Olszewski J, Lloyd-Smith D (1958). Focal seizures due to chronic localized encephalitis. *Neurology*, 8 (6), 435–445.

Swerdlow, J. L. (1995, June). Quiet miracles of the brain. *National Geographic*, 87, 2-41.

Vining, E. P., Freeman, J. M., Pillas, D. J., Uematsu, S., Carson, B. S., Brandt, J., Boatman, D., Pulsifer, M. B., & Zuckerman, A. (1997). Why would you remove half a brain? The outcome of 58 children after hemispherectomy-the Johns Hopkins experience: 1968 to 1996. *Pediatrics*, 100(2 Pt 1), 163-171.

find more hemispherectomy stories and information at: <http://hemifoundation.intuitwebsites.com/>

► Return to Lecture Guide

◄ Return to complete list of Lecture Launchers and Discussion Topics for Chapter 2

▲ Return to Chapter 2: Table of Contents

▼ CLASSROOM ACTIVITIES, DEMONSTRATIONS, AND EXERCISES

- 2.1 - Using Dominoes to Understand the Action Potential
- 2.2 - Stemming the Tide of Misinformation
- 2.3 - Environmental Influences on the Brain
- 2.4 - Demonstrating Neural Conduction: The Class as a Neural Network
- 2.5 - The Dollar Bill Drop
- 2.6 - Reaction Time and Neural Processing
- 2.7 - Mapping the Brain
- 2.8 - Football and Brain Damage
- 2.9 - Hemispheric Lateralization
- 2.10 - Hemispheric Communication and the Split Brain
- 2.11 - Crossword Puzzle
- 2.12 - Fill In the Blanks

[**▲ Return to Chapter 2: Table of Contents**](#)

Activity 2.1 - Using Dominoes to Understand the Action Potential

Walter Wagor suggests using real dominoes to demonstrate the so-called "domino effect" of the action potential as it travels along the axon. For this demonstration, you'll need a smooth table-top surface (at least 5 feet long) and one or two sets of dominoes. Set up the dominoes beforehand, on their ends and about an inch apart, so that you can push the first one over and cause the rest to fall in sequence. Proceed to knock down the first domino in the row and students should clearly see how the "action potential" is passed along the entire length of the axon. You can then point out the concept of refractory period by showing that, no matter how hard you push on the first domino, you will not be able to repeat the domino effect until you take the time to set the dominoes back up (i.e., the resetting time for the dominoes is analogous to the refractory period for neurons). You can then demonstrate the all-or-none characteristic of the axon by resetting the dominoes and by pushing so lightly on the first domino that it does not fall. Just as the force on the first domino has to be strong enough to knock it down before the rest of the dominoes will fall, the action potential must be there in order to perpetuate itself along the entire axon. Finally, you can demonstrate the advantage of the myelin sheath in axonal transmission. For this demonstration, you'll need to set up two rows of dominoes (approximately 3 or 4 feet long) next to each other. The second row of dominoes should have foot-long sticks (e.g., plastic rulers) placed end-to-end in sequence on top of the dominoes. By placing the all-domino row and the stick-domino row parallel to each other and pushing the first domino in each, you can demonstrate how much faster the action potential can travel if it can jump from node to node rather than having to be passed on sequentially, single domino by single domino. Ask your students to discuss how this effect relates to myelination.

Wagor, W. F. (1990). Using dominoes to help explain the action potential. In V. P. Makosky, C. C. Sileo, L. G. Whittemore, C. P. Landry, & M. L. Skutley (Eds.), *Activities handbook for the teaching of psychology: Vol. 3* (pp. 72-73). Washington, DC: American Psychological Association.

[**► Return to Lecture Guide**](#)

[**◄ Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2**](#)

[**▲ Return to Chapter 2: Table of Contents**](#)

Activity 2.2 - Stemming the Tide of Misinformation

Although there's a lot of promise in stem cell research, it comes with a lot of controversy as well. Consider these statements from various political figures over the years:

"While we must devote enormous energy to conquering disease, it is equally important that we pay attention to the moral concerns raised by the new frontier of human embryo stem cell research. Even the most noble ends do not justify any means."

GEORGE W. BUSH, speech, Aug. 9, 2001

"I think we can do ethically guided embryonic stem cell research. We have 100,000 to 200,000 embryos that are frozen in nitrogen today from fertility clinics. These weren't taken from abortion or something like that. They're from a fertility clinic, and they're either going to be destroyed or left frozen. And I believe if we have the option, which scientists tell us we do, of curing Parkinson's, curing diabetes, curing, you know, some kind of a ... you know, paraplegic or quadriplegic or, you know, a spinal cord injury — anything — that's the nature of the human spirit. I think it is respecting life to reach for that cure."

JOHN KERRY, presidential debate, Oct. 8, 2004

"The best that can be said about embryonic stem cell research is that it is scientific exploration into the potential benefits of killing human beings."

TOM DeLAY, *Washington Post*, May 25, 2005

"I am pro-life. I believe human life begins at conception. I also believe that embryonic stem cell research should be encouraged and supported."

BILL FRIST, speech, Jul. 29, 2005

"I'm very grateful that President Obama has lifted the restrictions on federal funding for embryonic stem cell research."

NANCY REAGAN, commentary, March 8, 2009

"The majority of Americans - from across the political spectrum, and of all backgrounds and beliefs - have come to a consensus that we should pursue this research. That the potential it offers is great, and with proper guidelines and strict oversight, the perils can be avoided."

BARACK OBAMA, executive order, March 9, 2009

Unfortunately, there's also a lot of misinformation about stem cells and stem cell research...in fact, one might question the scientific credentials of Mr. DeLay, whose noteworthy accomplishments (apart from a chequered political career) include running a pest control business and competing on *Dancing With the Stars*.

Encourage your students to examine the evidence and decide for themselves. Ask them to prepare a brief report on some aspect of stem cell research -- it's current legal status in the United States and worldwide; options for gathering stem cells; potential cures indicated by the current scientific evidence, and so on. You might ask different groups of students to tackle different issues, or ask all students to investigate a small set of issues. Similarly, you might structure this as a brief discussion exercise, or you might want to stage it as a more formal debate. There are many possible ways to implement this exercise. The important underlying aspect, however, is to get students to think critically and discuss openly the facts associated with stem cell research.

<http://stemcells.nih.gov/research/pages/current.aspx><http://www.tellmeaboutstemcells.org/>
<http://www.stemcells.com/view/0/index.html>
<http://www.physorg.com/news172072614.html>
http://www.wilsoncenter.org/index.cfm?event_id=161696&fuseaction=topics.event_summary&topic_id=116811
<http://www.biotech.ucdavis.edu/TBCWebsites/TBC07/StemCells&TissueEngineering/Li-MiraLoma/Biotech%20Website%20Design/index.html>
http://www.notable-quotes.com/s/stem_cells_quotes.html

► **Return to Lecture Guide**

◄ **Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2**

▲ **Return to Chapter 2: Table of Contents**

Activity 2.3 - Environmental Influences on the Brain

You might want to remind students that brain function and structure are subject to environmental influences. Ask students to identify the behaviors that are important for keeping the brain healthy and functioning well. The following are some possibilities:

Good nutrition, especially during childhood Adequate nutrition is vital for proper brain development. Even in adults, diet may influence brain function. Studies are showing that although high levels of cholesterol may be bad for your heart, low levels of cholesterol may be bad for the brain. Low cholesterol may be associated with low levels of the neurotransmitter serotonin, that can result in higher levels of aggression and depression.

Mental stimulation High levels of stimulation help to form neural connections that in turn enhance brain function.

Physical fitness Studies have shown that aerobic fitness has an impact on the density of capillaries in the brain. More capillaries result in greater blood flow to the brain.

Maternal health during pregnancy The uterine environment can have an enormous impact on the brain development of a fetus. Women who do not have adequate nutrition, or drink, smoke, or do drugs, and who are exposed to certain environmental toxins are more likely to have children with lower IQs and learning disabilities.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Activity 2.4 – Demonstrating Neural Conduction: The Class as a Neural Network

In this engaging exercise (suggested by Paul Rozin and John Jonides), students in the class simulate a neural network and get a valuable lesson in the speed of neural transmission. Depending on your class size, arrange 15 to 40 students so that each person can place his or her right hand on the right shoulder of the person in front of them. Note that students in every other row will have to face backwards in order to form a snaking chain so that all students (playing the role of individual neurons) are connected to each other. Explain to students that their task as a neural network is to send a neural impulse from one end of the room to the other. The first student in the chain will squeeze the shoulder of the next person, who, upon receiving this "message," will deliver (i.e., "fire") a squeeze to the next person's shoulder and so on, until the last person receives the message. Before starting the neural impulse, ask students (as "neurons") to label their parts; they typically have no trouble stating that their arms are axons, their fingers are axon terminals, and their shoulders are dendrites.

To start the conduction, the instructor should start the timer on a stopwatch while simultaneously squeezing the shoulder of the first student. The instructor should then keep time as the neural impulse travels around the room, stopping the timer when the last student/neuron yells out "stop." This process should be repeated once or twice until the time required to send the message stabilizes (i.e., students will be much slower the first time around as they adjust to the task). Next, explain to students that you want them to again send a neural impulse, but this time you want them to use their ankles as dendrites. That is, each student will "fire" by squeezing the ankle of the person in front of them. While students are busy shifting themselves into position for this exercise, ask them if they expect transmission by ankle-squeezing to be faster or slower than transmission by shoulder-squeezing. Most students will immediately recognize that the ankle-squeezing will take longer because of the greater distance the message (from

the ankle as opposed to the shoulder) has to travel to reach the brain. Repeat this transmission once or twice and verify that it indeed takes longer than the shoulder squeeze.

This exercise -- a student favorite -- is highly recommended because it is a great ice-breaker during the first few weeks of the semester, and it also makes the somewhat dry subject of neural processing come alive.

Rozin, P., & Jonides, J. (1977). Mass reaction time measurement of the speed of the nerve impulse and the duration of mental processes in class. *Teaching of Psychology*, 4, 91-94.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Activity 2.5 - The Dollar Bill Drop

After engaging in the neural network exercise, follow it up with the "dollar bill drop" (Fisher, 1979), which not only delights students but also clearly illustrates the speed of neural transmission. Ask students to get into pairs and to come up with one crisp, flat, one-dollar bill (or something larger, if they trust their fellow classmates!) between them. First, each member of the pair should take turns trying to catch the dollar bill with their nondominant (for most people, the left) hand as they drop it from their dominant (typically right) hand. To do this, they should hold the bill vertically so that the top, center of the bill is held by the thumb and middle finger of their dominant hand. Next, they should place the thumb and middle finger of their nondominant hand around the dead center of the bill, as close as they can get without touching it. When students drop the note from one hand, they should be able to easily catch it with the other before it falls to the ground.

Now that students are thoroughly unimpressed, ask them to replicate the drop, only this time one person should try to catch the bill (i.e., with the thumb and middle finger of the nondominant hand) while the other person drops it (i.e., from the top center of the bill). Student "droppers" are instructed to release the bill without warning, and "catchers" are warned not to grab before the bill is dropped. (Students should take turns playing dropper and catcher). There will be stunned looks all around as dollar bills whiz to the ground. Ask students to explain why it is so much harder to catch it from someone other than themselves. Most will instantly understand that when catching from ourselves, the brain can simultaneously signal us to release and catch the bill, but when trying to catch it from someone else, the signal to catch the bill can't be sent until the eyes (which see the drop) signal the brain to do so, which is unfortunately a little too late.

Fisher, J. (1979). *Body Magic*. Briarcliff Manor, NY: Stein and Day.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Activity 2.6 - Reaction Time and Neural Processing

Yet another exercise that illustrates the speed of neural processing is suggested by E. Rae Harcum. The point made by this simple but effective exercise is that reaction times increase as more response choices become available (i.e., because more difficult choices in responses involve more neuronal paths and more synapses, both of which slow neural transmission). Depending on your class size, recruit two equal groups of students (10 to 20 per group is ideal) and have each group stand together at the front of the room. First, explain that all subjects are to respond as quickly as possible to the name of a U.S. President. Then give written instructions to each group so that neither group knows the instructions given to the other. One group should be instructed to raise their right hands if the president served before Abraham Lincoln and to raise their left hands if the president served after Lincoln. The other group should be instructed simply to raise their left hands when they hear a president's name. Ask participants and

Copyright © 2017 Pearson Education. All rights reserved.

audience members to note which group reacts more quickly. When all students are poised and ready to go (i.e., hands level with shoulders and ready to raise), say "Ready" and then "Reagan." The group with the simpler reaction time task should be faster than the group whose task requires a choice.

Harcum, E. R. (1988). Reaction time as a behavioral demonstration of neural mechanisms for a large introductory psychology class. *Teaching of Psychology*, 4, 208-209.

► [Return to Lecture Guide](#)


◄ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Activity 2.7 - Mapping the Brain

Many students, especially those with little background in the sciences, will find it a challenge to keep track of the location of all the parts of the brain outlined in the text. One simple way to reinforce their learning of brain structure is to have students locate the various parts on a diagram of the brain. The brain diagram and the student instructions for this exercise are included as **Handout Master 2.1**. The day before you present this activity, ask students to bring colored pencils or markers to the next class meeting. On the day of the activity, divide students into small groups and distribute copies of the diagram of the brain and the accompanying questions in the student handouts. Within their groups students can help each other locate each part of the brain and then color code them using their pencils or markers. They can also indicate the function of each part on the diagram. This exercise is very useful for helping students to memorize brain anatomy, and the color-coded diagram serves as a helpful study guide.

For your convenience, a completed diagram and suggested answers to the questions are furnished here.



1. This is a diagram of the left side of the brain.

Left side functions: The left hemisphere controls touch and movement of the right side of the body, vision in the right half of the visual field, comprehension and production of speech, reading ability, mathematical reasoning, and a host of other abilities.

Right side functions: The right hemisphere controls touch and movement of the left side of the body,

Copyright © 2017 Pearson Education. All rights reserved.

vision in the left half of the visual field, visual-spatial ability, map-reading, art and music appreciation, analysis of nonverbal sounds, and a host of other abilities.

2. The front of the brain is on the left side of the diagram; the back of the brain is on the right.
3. The cerebrum is the sum of the frontal, parietal, temporal, and occipital lobes. The cerebellum is labeled on the diagram above. The cerebrum is responsible for higher forms of thinking, including a variety of specific abilities described under motor cortex, visual cortex, somatosensory cortex, and auditory cortex. The cerebral cortex also contains vast association areas, whose specific functions are poorly defined but may include reasoning and decision making, planning appropriate behavior sequences, and knowing when to stop. The limbic system, which appears to be strongly involved in regulating emotions, is also part of the cerebrum. The cerebellum aids in the sense of balance and motor coordination.
4. The frontal, parietal, temporal, and occipital lobes are labeled on the diagram above.
5. The motor cortex is labeled on the diagram above. The motor cortex in each hemisphere controls movements on the opposite side of the body.
6. The visual cortex is labeled on the diagram above. The visual cortex in each hemisphere receives information from the visual field on the opposite side.
7. The auditory cortex is labeled on the diagram above. The auditory cortex is responsible for processing sounds.
8. The somatosensory cortex is labeled on the diagram above. The somatosensory cortex on each side receives information about touch, joint position, pressure, pain, and temperature from the opposite side of the body.
9. Broca's and Wernicke's areas are labeled on the diagram above. Broca's area is often referred to as the motor speech area. It is responsible for our ability to carry out the movements necessary to produce speech. Wernicke's area is often referred to a sensory speech area. It is mainly involved in comprehension and planning of speech.
10. Neurons would be found all over the drawing. (The brain is made up of billions of neurons.) Each neuron is very tiny compared to the size of the brain, so no single neuron would be visible to the naked eye in a drawing at this scale. The cell bodies of the largest neurons in the brain are about 1/20 of a millimeter in diameter!
11. The brain stem is labeled on the diagram above. Different parts of the brain stem are involved in regulation of sleep and wakefulness, dreaming, breathing, heart rate, and attentional processes.

► **Return to Lecture Guide**

◄ **Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2**

▲ **Return to Chapter 2: Table of Contents**

Activity 2.8 - Football and Brain Damage

Coaches and medical experts have known for a while that the severe hits that football players take on the field can lead to concussions, blacking out, and even permanent damage. More recently, however, there has been increasing concern that the effects of repeated hits to the head may not manifest themselves until decades later. Early studies suggest that former NFL players suffer high rates of memory and other cognitive problems years after retiring, and that they also may develop these problems earlier than non-football players do. NFL players may also be vulnerable to higher rates of depression and Alzheimer's disease.

To investigate this problem, groups like the Sports Legacy Institute have begun to encourage former NFL players to donate their brains to science when they die. Already, the brains of a handful of players have been examined, with shocking results. Almost all of the brains show high levels of a protein called *tau*, which is suspected of being involved in several neurodegenerative disorders, including Alzheimer's disease. The presence of high levels of tau may explain why football players have a tendency to develop cognitive impairments long after their playing days are over. More disturbing still, high levels of tau have also been found in the brain of an 18 year-old high school football player who died.

After introducing students to this issue, have the class discuss the possible implications for social and sports policy. Should football playing be stopped? Should the rules of the game be changed to eliminate hard hitting? If necessary, pose the following additional questions to stimulate discussion: Everyone knows football is dangerous, but does the fact that these cognitive impairments may take decades to develop make them somehow different? Is the risk of permanent cognitive disability somehow different than the risk of permanent physical disability? Wrestlers, soccer players, boxers, and other types of athletes are also at risk for long term brain damage. Should these sports be changed or banned?

After discussing the issue in class, have students respond to the following writing prompt.

Writing Prompt: Describe a longitudinal and then a cross-sectional study that could be used to determine if professional football players show higher than normal rates of cognitive impairment. Explain some of the advantages and disadvantages of the two designs.

Sample answer: *A longitudinal study might choose a few football players and then test them every 10 years using the same cognitive tests to see how their abilities change over time. A cross-sectional study, on the other hand, might find a group of 65 year old retired football players and compare their cognitive functioning to 65 year olds who did not play football. The longitudinal study would provide a more complete view of how cognitive function might decline, but would take decades to complete, and may suffer from attrition. The cross-sectional study would be a lot easier to perform, but would only offer a "snap-shot" of cognitive function. You could not tell, for example, if football players develop cognitive impairment earlier than non-football players typically do.*

Miller, G. (2009). A late hit for pro football players. *Science*, 7, 670–672.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Activity 2.9 - Hemispheric Lateralization

Hemispheric lateralization results in eyedness, handedness, footedness, earedness, facedness, and other silly-sounding words with important implications (see the related Lecture Launcher in this chapter). Lateralization results from the specialization of each hemisphere for different tasks, such as reading facial expressions, speaking, solving spatial problems, or performing analytic tasks. Although neuropsychologists use sophisticated measures to determine these lateralization, this simple exercise allows students to gauge their own brain organization.

With both eyes open, have students hold up their right thumbs at arm's length under an object across the room directly in front of them. As they alternately close their left and right eyes, their thumbs should appear to jump to the right or to the left with respect to the distant object. For those who are right-eyed, their thumbs should jump to the right when they close their right eyes, but stay as is when they close their left eyes. The opposite pattern should occur among those who are left-eyed. Students who see little or no jumping are among the 41% of the population who are neither strongly left-eyed nor right-eyed.

As a second test, ask for a volunteer. Present the student with the first paragraph of this exercise (or any suitable short passage) to memorize, a broom, a clock with a second hand, a pencil, and a pad of paper.

First, time how long the volunteer can balance the broom on the tip of his or her right index finger while standing on the right foot. Next, measure the time as the volunteer balances the broom on his or her left index finger while standing on the left foot. Finally, repeat these tests while the volunteer recites the memorized passage. Speech will be localized on the side of the brain opposite the hand that is most disrupted by the memorization task.

Another demonstration, suggested by Morton Ann Gernsbacher, requires students to move their right hand and right foot simultaneously in a clockwise direction for a few seconds. Next ask that the right hand and left foot be moved in a clockwise direction. Then, have students make circular movements in opposite directions with right the hand and the left foot. Finally, have students attempt to move the right hand and right foot in opposite directions. This generally produces laughter as students discover that this procedure is most difficult to do even though they are sure – before they try it – that it would be no problem to perform. A simple alternative activity is to ask students to pat their heads and to rub their stomachs clockwise and then switch to a counterclockwise motion. The pat will show slight signs of rotation as well.

The brain is lateralized to some extent, and this makes some activities difficult to perform. Challenge your students to explain why activities of these types are difficult to execute. This will generally lead to interesting discussions and the assertion by some students that this type of behavior is no problem. Generally students who have been trained in martial arts, dance and/or gymnastics have less difficulty completing these activities due to their rigorous physical training.

Haseltine, E. (1999, June). Brain works: Your better half. *Discover*, 112.

Kemle, E. D. (1987). Cerebral lateralization. In V. P. Makosky, L. G. Whittemore, and A. M. Rogers (Eds.). *Activities handbook for the teaching of psychology* (Vol. 2) (pp. 33–36). Washington, D.C.: American Psychological Association.

Kemle, E. D., Filipi, T., & Gravlin, L. (1985). Some simple classroom experiments on cerebral lateralization. *Teaching of Psychology*, 12, 81–83.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Activity 2.10 - Hemispheric Communication and the Split Brain

Even after reading the textbook and listening to your lecture, many students may have difficulty conceptualizing the effects of a split-brain operation on an individual's behavior. Morris (1991) described five activities designed to simulate the behavior of split-brain patients. All of the activities have the same basic setup. You will need to solicit two right-handed volunteers and seat them next to each other at a table, preferably in the same chair. The volunteer on the left represents the left hemisphere, and the other student is the right hemisphere. The students are instructed to place their outer hand behind their back and their inner hands on the table with their hands crossed, representing the right and left hands of the split-brain patient. Finally, the student representing the right hemisphere is instructed to remain silent for the remainder of the activity. In one of the activities described by Morris, both students are blindfolded and a familiar object (Morris suggested a retractable ball-point pen) is placed in the left hand of the "split-brain patient" (the hand associated with the right hemisphere). Then ask the "right hemisphere" student if he or she can identify the object, reminding him or her that they must do so nonverbally. Next, ask the "right hemisphere" to try to communicate, without using language, what the object is to the "left hemisphere." Your more creative volunteers may engage in behaviors that attempt to communicate what the object is through sound or touch. If your "right hemisphere" has difficulty in figuring out how to communicate, ask the class for suggestions. This demonstration can be used to elicit discussion about why only the "left hemisphere" student can talk, the laterality of the different senses, and how split-brain patients are able to adjust their behavior to accommodate. You should refer to Morris's original article for descriptions of the other activities.

Morris, E. J. (1991). Classroom demonstration of behavioral effects of the split-brain operation. *Teaching of Psychology*, 18, 226-228.

► [Return to Lecture Guide](#)

◄ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Activity 2.11 - Crossword Puzzle

Copy and distribute [Handout Master 2.2](#) to students as a homework or in-class review assignment.

Answers for the Crossword puzzle:

Across

1. neurotransmitter that causes the receiving cell to stop firing. **Inhibitory**
3. the cell body of the neuron, responsible for maintaining the life of the cell. **Soma**
4. endocrine gland located near the base of the cerebrum which secretes melatonin. **Pineal**
7. glands that secrete chemicals called hormones directly into the bloodstream. **Endocrine**
8. long tube-like structure that carries the neural message to other cells. **Axon**
10. chemical found in the synaptic vesicles which, when released, has an effect on the next cell.

Neurotransmitter

13. bundles of axons coated in myelin that travels together through the body. **nerves**
14. branch-like structures that receive messages from other neurons. **Dendrites**
15. endocrine gland found in the neck that regulates metabolism. **Thyroid**
17. thick band of neurons that connects the right and left cerebral hemispheres. **Corpus Callosum**
19. part of the nervous system consisting of the brain and spinal cord. **Central**

Down

2. part of the limbic system located in the center of the brain, it acts as a relay from the lower part of the brain to the proper areas of the cortex. **Thalamus**
4. endocrine gland that controls the levels of sugar in the blood. **Pancreas**
5. fatty substances produced by certain glial cells that coat the axons of neurons to insulate, protect, and speed up the neural impulse. **Myelin**
6. the basic cell that makes up the nervous system and which receives and sends messages within that system. **Neuron**
8. chemical substances that mimic or enhance the effects of a neurotransmitter on the receptor sites of the next cell. **Agonists**
9. part of the lower brain that controls and coordinates involuntary, rapid, fine motor movement.

Cerebellum

11. process by which neurotransmitters are taken back into the synaptic vesicles. **Reuptake**
12. a group of several brain structures located under the cortex and involved in learning, emotion, memory, and motivation. **Limbic**
16. chemicals released into the bloodstream by endocrine glands. **Hormones**
18. brain structure located near the hippocampus, responsible for fear responses and memory of fear.

Amygdala

► [Return to Lecture Guide](#)

◄ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Activity 2.12 - Fill-in-the-Blanks

Copy and distribute [Handout Master 2.3](#) to students as a homework or in-class review assignment.

Answers for Fill-in-the-Blanks—Chapter 2

nervous system
neuron

axon
dendrites
soma
myelin
nerves
ions
resting potential
synaptic vesicles
neurotransmitters
excitatory
agonists
spinal cord
sensory
peripheral nervous
somatic nervous
autonomic nervous
sympathetic division
electroencephalograph
cerebellum
thalamus
pons
reticular formation
hippocampus
amygdala
cortex
corpus callosum
occipital cortex
parietal cortex
temporal lobes
frontal lobes
endocrine
adrenal glands

► Return to Lecture Guide

◄ Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2

▲ Return to Chapter 2: Table of Contents

▼ HANDOUT MASTERS

- > [Handout Master 2.1 Mapping the Brain](#)
- > [Handout Master 2.2 Crossword Puzzle](#)
- > [Handout Master 2.3 Fill-in-the-Blanks](#)

◀ [Return to complete list of Classroom Activities, Demonstrations, and Exercises for Chapter 2](#)

▲ [Return to Chapter 2: Table of Contents](#)

Handout Master 2.1
Mapping the Brain

Label the diagram of the brain to show or answer the following questions.

1. Is this a drawing of the left side or the right side of the brain? What are the particular functions of that side of the brain as compared to the other hemisphere?

Left side functions:

Right side functions:


2. Where is the front of the brain? Where is the back?
3. Label the cerebrum and cerebellum and describe their functions.

Cerebral functions:

Cerebellar functions:


4. Label the four lobes of the cerebral cortex.
5. Label the motor cortex and describe its function.
6. Label the visual cortex and describe its function.
7. Label the auditory cortex and describe its function.
8. Label the somatosensory cortex and describe its function.
9. Label Broca's and Wernicke's areas and describe their functions.
10. Where would you expect to find neurons in this drawing and how big would they be if they were drawn?
11. Label the brain stem. What is its function?

► Return to Activity: Mapping the Brain
◄ Return to complete list of Handout Masters for Chapter 2
▲ Return to Chapter 2: Table of Contents



Handout Master 2.2

Crossword Puzzle



Across

1. Neurotransmitter that causes the receiving cell to stop firing.
3. The cell body of the neuron, responsible for maintaining the life of the cell.
4. Endocrine gland located near the base of the cerebrum which secretes melatonin.
7. Glands that secrete chemicals called hormones directly into the bloodstream.
8. Long tube-like structure that carries the neural message to other cells.
10. Chemical found in the synaptic vesicles which, when released, has an effect on the next cell.
13. Bundles of axons coated in myelin that travel together through the body.
14. Branch-like structures that receive messages from other neurons.
15. Endocrine gland found in the neck that regulates metabolism.
17. Thick band of neurons that connects the right and left cerebral hemispheres.
19. Part of the nervous system consisting of the brain and spinal cord.

Down

2. Part of the limbic system located in the center of the brain, it acts as a relay from the lower part of the brain to the proper areas of the cortex.
4. Endocrine gland that controls the levels of sugar in the blood.
5. Fatty substances produced by certain glial cells that coat the axons of neurons to insulate, protect, and speed up the neural impulse.
6. The basic cell that makes up the nervous system and which receives and sends messages within that system.
8. Chemical substances that mimic or enhance the effects of a neurotransmitter on the receptor sites of the next cell.
9. Part of the lower brain that controls and coordinates involuntary, rapid, fine motor movement.
11. Process by which neurotransmitters are taken back into the synaptic vesicles.
12. A group of several brain structures located under the cortex and involved in learning, emotion, memory, and motivation.
16. Chemicals released into the bloodstream by endocrine glands.
18. Brain structure located near the hippocampus, responsible for fear responses and memory of fear.

► Return to Activity: Crossword Puzzle

◄ Return to complete list of Handout Masters for Chapter 2

▲ Return to Chapter 2: Table of Contents

Handout Master 2.3
Fill-in-the-Blanks

1. An extensive network of specialized cells that carry information to and from all parts of the body is called the _____.
2. The basic cell that makes up the nervous system and which receives and sends messages within that system is called a _____.
3. The long tube-like structure that carries the neural message to other cells on the neuron is the _____.
4. On a neuron, the branch-like structures that receive messages from other neurons is the _____.
5. The cell body of the neuron, responsible for maintaining the life of the cell and contains the mitochondria is the _____.
6. The fatty substances produced by certain glial cells that coat the axons of neurons to insulate, protect, and speed up the neural impulse is the _____.
7. The bundles of axons in the body that travel together through the body are known as the _____.
8. The charged particles located inside and outside of the neuron are called _____.
9. The state of the neuron when not firing a neural impulse is known as the _____.
10. The _____ are sack-like structures found inside the synaptic knob containing chemicals.
11. _____ are chemicals found in the synaptic vesicles which, when released, has an effect on the next cell.
12. The _____ neurotransmitter causes the receiving cell to fire.
13. The _____ mimic or enhance the effects of a neurotransmitter on the receptor sites of the next cell, increasing or decreasing the activity of that cell.
14. The _____ a long bundle of neurons that carries messages to and from the body to the brain that is responsible for very fast, lifesaving reflexes.
15. A neuron that carries information from the senses to the central nervous system and is also known as the afferent is called a _____.
16. All nerves and neurons that are not contained in the brain and spinal cord but that run through the body itself are in the _____ system.
17. The division of the PNS consisting of nerves that carry information from the senses to the CNS and from the CNS to the voluntary muscles of the body is the _____ system.
18. The _____ system division of the PNS consisting of nerves that control all of the *involuntary* muscles, organs, and glands sensory pathway nerves coming from the sensory organs to the CNS consisting of sensory neurons.
19. The part of the ANS that is responsible for reacting to stressful events and bodily arousal is called the _____ of the nervous system.
20. A machine designed to record the brain wave patterns produced by electrical activity of the surface of the brain is called an _____.
21. The part of the lower brain located behind the pons that controls and coordinates involuntary, rapid, fine motor movement is called the _____.
22. The part of the limbic system located in the center of the brain, this structure relays sensory information from the lower part of the brain to the proper areas of the cortex and processes some sensory information before sending it to its proper area and is called the _____.

23. The larger swelling above the medulla that connects the top of the brain to the bottom and that plays a part in sleep, dreaming, left–right body coordination, and arousal is called the _____.
24. The _____ is an area of neurons running through the middle of the medulla and the pons and slightly beyond that is responsible for selective attention.
25. The _____ is a curved structure located within each temporal lobe, responsible for the formation of long-term memories and the storage of memory for location of objects.
26. The _____ is a brain structure located near the hippocampus, responsible for fear responses and memory of fear.
27. The _____ is the outermost covering of the brain consisting of densely packed neurons, responsible for higher thought processes and interpretation of sensory input.
28. The thick band of neurons that connects the right and left cerebral hemispheres is called the _____.
29. The section of the brain located at the rear and bottom of each cerebral hemisphere containing the visual centers of the brain is the called the _____.
30. The sections of the brain located at the top and back of each cerebral hemisphere containing the centers for touch, taste, and temperature sensations is called the _____.
31. The _____ is the area of the cortex located just behind the temples containing the neurons responsible for the sense of hearing and meaningful speech.
32. The _____ are areas of the cortex located in the front and top of the brain, responsible for higher mental processes and decision making as well as the production of fluent speech.
33. The _____ glands secrete chemicals called hormones directly into the bloodstream.
34. The endocrine glands located on top of each kidney that secrete over 30 different hormones to deal with stress, regulate salt intake, and provide a secondary source of sex hormones affecting the sexual changes that occur during adolescence are called the _____.

Words for Fill-in-the Blanks

adrenal glands
agonists
amygdala
autonomic nervous
axon
cerebellum
corpus callosum
cortex
dendrites
electroencephalograph
endocrine
excitatory
frontal lobes
hippocampus
ions
myelin
nerves
nervous system
neuron
neurotransmitters
occipital cortex
parietal cortex
peripheral nervous
pons
resting potential
reticular formation
sensory
soma
somatic nervous
spinal cord
sympathetic division
synaptic vesicles
temporal lobes
thalamus

► **Return to Activity: Fill-in-the-Blanks**

◄ **Return to complete list of Handout Masters for Chapter 2**

▲ **Return to Chapter 2: Table of Contents**

▼ VIDEO SERIES

The VIDEO SERIES features over 100 original videos covering the most recent research, science, and applications across the general psychology curriculum, and utilizing the latest in film and animation technology. Each 4-6 minute video clip has automatically graded assessment questions tied to it.

Access www.MyPsychLab.com, and select “Videos.”

Chapter 2 Video Content available:

Video: The Basics: How the Brain Works, Part 1 (5:04)

Find out about neurons: their parts, how they communicate, and their function in the brain.

Video: The Basics: How the Brain Works, Part 2 (6:11)

Find out about the nervous system: how it's divided and processes information, the significance of each brain structure, and how neuronal transmission works.

Video: The Big Picture: My Brain Made Me Do It. (4:59)

Learn about the function of different parts of the nervous system and how it integrates multiple signals to create experience.

Video: In the Real World: Neurotransmitters (3:10)

Learn about the neurotransmitter dopamine and what happens when there is an imbalance of dopamine as a result of natural causes or drugs.

Video: Special Topics: The Plastic Brain (7:04)

Learn specifics about how the human brain grows and develops throughout the lifespan and how the brain can remarkably repair itself or compensate after being damaged.

Video: Thinking Like a Psychologist: The Pre-Frontal Cortex (3:28)

See how certain areas of the brain contribute to different behaviors, and how both biological and environmental factors can lead a person to become a violent criminal later in life.

Video: What's In It For Me? Your Brain on Drugs (3:51)


Neurotransmitter imbalances can occur from chemicals that we willingly ingest, such as drugs or alcohol.

▲ Return to Chapter 2: Table of Contents

▼ VISUAL BRAIN

THE VISUAL BRAIN is an interactive virtual brain designed to help students better understand neuroanatomy, physiology, and human behavior. Thirteen virtual brain modules bring to life many of the most difficult topics typically covered in the introductory course. This hands-on experience engages students and helps make course content and terminology relevant. Modules relevant to the current chapter are highlighted in **bold** below.

Access www.MyPsychLab.com, and click on the Student Resources tab.



Select among these modules

- **Brain Damage and Neuroplasticity**
- **Control of Movement**
- Drug Addiction
- Hunger and Eating
- **Lateralization and Language**
- Learning and Memory
- **Nervous System**
- **Neural Conduction**
- Perception
- Psychiatric Disorders
- Sleep and Waking
- Stress and Health
- **Visual System**

▲ Return to Chapter 2: Table of Contents

▼ EXPERIMENT AND SURVEY SIMULATIONS

EXPERIMENT AND SURVEY SIMULATIONS allow students to participate in online simulations of virtual versions of classic psychology experiments, surveys, and research-based inventories, helping to reinforce what they are learning in class and from their textbook.

Access www.MyPsychLab.com, and select “Simulations.”

Chapter 2 Simulations Content available:

Simulation: Hemispheric Specialization

Judge whether a string of letters in your peripheral vision is a real word and test whether language functions are lateralized in the brain.

Survey: Do You Fly or Fight?

Participate in a survey to discover if you flee or fight under stressful conditions.

▲ Return to Chapter 2: Table of Contents

▼ **WRITING SPACE** *featuring* auto-feedback **WRITING PRACTICE**

WRITING SPACE helps students master concepts and develop critical thinking through writing. WRITING SPACE provides a single place within your MyLab to create and track your writing assignments, access writing resources, and exchange meaningful, personalized feedback quickly and easily. Plus, WRITING SPACE has integrated access to Turnitin, the global leader in plagiarism prevention.

Access www.MyPsychLab.com, and select “Writing Space” from the left-hand navigation bar.

Auto-feedback Writing Practice accessible from Writing Space

WRITING PRACTICE prompts within WRITING SPACE offer immediate automated feedback. Each student submission receives detailed feedback based on the following traits: Development of Ideas, Organization, Conventions, Voice, Focus and Coherence. Instructors can provide additional feedback and can adjust the auto-generated grade.

WRITING PRACTICE SAMPLE 2.1

Your best friends invite you over for pizza and a friendly game of cards. Describe how the following parts of the brain are involved during your evening of eating pizza, socializing, and playing cards: Broca's area, hippocampus, hypothalamus, and occipital lobe.

Instructor-Created Writing Assignments accessible from Writing Space:

Instructors can create their own writing prompts and grading rubrics, or copy and paste from a library of sample prompts and rubrics available within this Instructor Manual. Instructors can provide personalized feedback and grades to students.

INSTRUCTOR-CREATED SAMPLE 2.1

The news media sometimes report functional brain imaging findings accurately, but often report them in oversimplified ways, such as implying that researchers have identified a single brain region for Capacity X (like religious beliefs, morality, or political affiliation). Locate two media reports on functional brain imaging (ideally using fMRI or PET) and evaluate the quality of media coverage. Did the reporters interpret the findings correctly, or did they go beyond the findings? For example, did the reporters avoid implying that investigators located a single brain “spot” or “region” underlying a complex psychological capacity?

▲ Return to Chapter 2: Table of Contents